

العدد الثاني عشر - ديسمبر 2016

Micro- scale preparation of isotopically enriched monomethylmercury (MM²⁰¹Hg)

Abdelkarem A. Elgazali^{1,2}, Yayoi Kobayashi³, Eva M. Krupp⁴ and Jörg Feldmann⁴

¹ Chemistry Department, Faculty of Arts and Science, University of Benghazi, Al-Marj city, Libya,

² Environmental and Biological Chemistry research Centre, Tukra, University Of Benghazi, Libya

³ Environmental Health Sciences Division, National Institute for Environmental Studies, Japan

⁴ Chemistry department, University of Aberdeen, Aberdeen, Scotland, UK, AB24 3UE



Abstract

A simple and fast synthesis method for the micro-scale laboratory preparation of isotopically enriched monomethylmercury (MMHg) which is not commercially available has been evaluated and established successfully. This enriched solution is an important standard for species-specific isotope dilution mass spectrometry analysis (SSIDMS) for determination of the true concentration of methylmercury in biological (hair) and environmental samples (soil and sediment). The isotopically enriched MMHg has been synthesised from inorganic standard solution prepared from commercially available mercury oxide (^{201}HgO) using methylcobalamin co-enzyme as methylation agent. The synthesis time required is 1 h at 37°C . The product is highly pure, yielded more than 95% as ^{201}Hg in MMHg. The proposed method allows control of unintentional formation of health hazard dimethylmercury. It also allows work on a micro-scale to control the use of expensive enriched isotope standard. The synthesised product was analysed using capillary gas chromatography (CGC) coupled with inductively coupled plasma mass spectrometry (ICP-MS) after derivatisation with sodium tetraethylborate (NaBPr_4). The concentration, purity, isotopic composition and stability of synthesised, enriched MMHg have been investigated in order to establish standard protocols for MMHg isotope dilution analysis.

Keywords: Monomethylmercury; species-specific isotope dilution mass spectrometry analysis (SSIDMS); Sodium tetra (n-propyl) borate; mercury oxide (^{201}HgO); Dimethylmercury; methylcobalamin co-enzyme.

Introduction:

Mercury is potentially one of the most toxic metals to organisms as it has strong ability to bioaccumulate and the enrichment of highly toxic mercury (Hg) compounds in aquatic food chain poses a serious environmental problem (1-5). The most important chemical forms of mercury are elemental mercury (Hg^0), inorganic mercury (Hg^{2+}), monomethylmercury (Me-Hg, CH_3Hg^+) and dimethylmercury (DMHg, CH_3HgCH_3). In the biogeochemical cycle of mercury, the mercury vapour is released to the environment through anthropogenic sources i.e. human activity including emissions from coal burning in power station, chloro-alkali plants and artisanal gold mining which is oxidized in the upper atmosphere to water-soluble ionic mercury (Hg^{++}), and returned to earth surface (both soil and water) in rainwater(1,6).

The long transport cycle of mercury in the atmosphere; its deposition, bioaccumulation, and the enrichment of highly toxic methylmercury (Me-Hg) compounds in aquatic food chain pose a serious environmental problem, even in remote areas and contaminating, humans(3,4,7-9). MMHg is efficiently adsorbed from the gastro-intestinal tract, and it passes the blood-brain and placenta barriers and it causes disintegration of cells within the brain. The main target of methylmercury in humans is the central nervous system- especially the control sensory, visual and auditory areas involved in coordination. The most severe effects cause widespread brain damage, resulting mental confusion, blindness, coma and death (10-13). On the other hand, the most common forms of exposure to inorganic mercury forms are by inhalation of mercury (Hg) vapour released from chlor alkali plants, dental amalgam and gold mining activities. During the ingestion of inorganic mercury at high concentration, the corrosive effects first of all, damages the digestive tract, causing vomiting and stomach pain, and in severe cases, may result in shock and finally, renal tubule degeneration, and kidneys dysfunction may be seen (14). The exposure of inorganic compounds (In-Hg) (1-5% contents) may cause irritation, contact dermatitis and corrosion of the skin (15).

Regulatory authorities are required to measure mercury (Hg) in a variety of food, industrial, environmental and biological samples for reasons of human being health. In addition to total mercury (T-Hg) measurement in the environmental, food and biological samples, the specific measurement of methylmercury (Me-Hg), noted Me-Hg is of concern to regulatory organisation; as this Hg species has a mammalian LD50 1000 times lower than elemental mercury. Moreover, many foodstuffs such as fish contain the majority of mercury (Hg) as methylmercury (Me-Hg). In addition based on the results of provisional tolerable weekly intakes (PTWIs) from Seychelles, New Zealand and Faroe Island studies, in year 2000 the United States Environmental Protection Agency (U.S. EPA) developed a reference Doses (RfD) for methylmercury of $0.1\mu g$. Me-Hg/kg body weight/day, which were equivalent to a hair mercury concentration of $1.0\mu g$. Hg/ g (15,16).

In the year 2003, the JECFA, the joint FAO/WHO (Food and Agriculture Organisation of the United Nations and World Health Organisation) Expert Committee on Food Additives recommended that the provisionally human consumption of Me-Hg is limited to $1.6\mu g$ per kg body mass, per week that is equivalent to a hair mercury concentration of about $2.3\mu g$. Hg/g hair(15,16). Therefore, it is essential to measure the exact concentration of methylmercury and inorganic mercury present in biological, environmental and food samples through accurate analytical methods to be used in control laboratories.

العدد الثاني عشر - ديسمبر 2016

Analytical techniques for separation of methylmercury (Me-Hg) are well documented(17). After extraction from solid matrices and derivatization, the methylmercury (Me-Hg) is frequently measured using hyphenated techniques(17). Mercury speciation analysis has been usually performed by resorting to hyphenated techniques, based on the coupling of an effective separation technique to a sensitive element-specific detector. Capillary gas chromatography (CGC), liquid chromatography (LC) or more recently capillary electrophoresis (EC) can be interfaced with specific atomic detection including atomic absorption spectrometry (AAS), atomic fluorescence spectrometry (AFS), electron capture, or inductively coupled plasma mass spectrometry (ICPMS) (10,18-24). recently researches has shown that the coupling of GC to ICP-MS appear to be more suitable hyphenated technique to carry out the mercury speciation analysis because of its high sensitivity, multi isotopic, and multi elemental capabilities(17,18,22). However, the GC-ICP-MS is only suitable for volatile species like mercury species(19,22).

Quality results are sometimes associated with sample pre-treatment, for the analysis of solids such as biological and environmental samples required a leaching (alkaline or acid) /digestion step to liberate mercury species from sample matrix prior to detection with GC-ICP-MS. However, for ionic mercury species derivatization reaction are required in order to achieve good results. Moreover, there are potential sources of error related to non-quantitative recoveries of mercury species, artefact formation and methylmercury transformation during solid matrixes sample preparation and derivatisation steps (19,25). For example, Liang and Lazoff reported that the methylmercury artefacts were formed in alkaline digestion step for sediment samples ranging from 0.001 to 0.002% of the total mercury (T-Hg) (spiked plus un-spiked) concentration (26).

To tackle these potential problems of speciation data validation in order to calculate the original level of methylmercury, the use of isotope dilution mass spectrometry techniques offers great potential of very small uncertainties since quantitative recoveries are not required and rearrangement reactions are easily detected(10,24,27). Provided that the enriched isotope can perform the role of ideal internal standard since it is present in an equilibrated and equivalent state to the natural isotope (28). Isotope dilution mass spectrometry technique (IDMS) has been considered to be a best analytical method for speciation and is a well known based on the measurement of isotope ratio in samples where its isotopic composition has been altered by the addition of a known amount of an isotopically enriched element(17,23,29). A main advantage of this technique (IDMS) that chemical separation, if required for accurate ratio determination, need not be quantitative. Moreover, concentrations of chemical species can be measured very precisely because ratios can be measured very reproducibly(17,29).

However, in spite of the benefits of Isotope dilution mass spectrometry technique (IDMS), it is not being used widely as a method of analysis because the lack of suitable enriched internal calibration standards. Due to the isotopically enriched methylmercury standard is not commercially available therefore needs to be synthesized in the laboratory.

According to the several literature surveys, there are some proposed methods for the production of organomercury compounds, e.g. the reaction between dimethylmercury (DMHg) and inorganic mercury ($HgCl_2$) the chemically methylate of mercury (II) by methylcobalamin in the absent of cell extract and the reaction of tetramethyltin ($(CH_3)_4Sn$) with mercuric chloride ($HgCl_2$) stock solutions(10,28). However, the most often used method for production of organomercury compounds is based on the reaction of methylcobalamin

العدد الثاني عشر - ديسمبر 2016

(CH₃ CO, a vitamin B12 analog) with inorganic mercury (In-Hg). In this method, there are two reaction products; monomethylmercury (MMHg) chloride and dimethylmercury (DMHg) chloride were produced with different yield depends mainly on the reaction time, temperature and the mole ratio of the reactants used(10). The major focus of most of these studies was the reaction product of the methylcobalamin ortetramethyltin ((CH₃)₄Sn) with inorganic mercury (In-Hg), not the synthesis of enriched methylmercury with high yield and impurity in order to use it as a standard compound. However, only few methods for synthesis of isotopically enriched monomethylmercury chloride have been found in the literature.

Rouleau C and Block M (30) synthesized the radioactive methylmercury chloride (CH₃²⁰³Hg(II)) by methylation of inorganic mercury (²⁰³Hg (II)) with methylcobalamin (MeCo) and isolation of CH₃²⁰³Hg(II) from the reaction mixture in a single extraction step with hexane / benzene (1:1) and the combined organic layers were stirred over sodium carbonate (NaCO₃) solution. The synthesized the radioactive methylmercury chloride (CH₃²⁰³Hg (II)) was left dissolved in (NaCO₃) solution in a suitable. The final yield was ≥90% and the time required was less than 4 hours.

On the other hand R. C. Rodriguez Martin-Domimeadiset al.(28) synthesized the isotopically enriched monomethylmercury (MMe²⁰¹Hg) by methylation of commercially available mercury oxide (²⁰¹HgO) using methylcobalamin co-enzyme as methylation reagent. The mixture was left sitting in the dark place at 37°C in water bath for 1 hour. In order to stop the methylating reaction and to convert the unintentionally formed DMeHg into methylmercury, 1 ml of HCl (conc.) was added to the solution at cooled at 4°C and then the solution was shaken for 5 min. Finally the micro-scale synthesized of isotopically enriched MMe²⁰¹Hg was extracted three times with 500 ul of toluene and the combination extracts were dried over sodium sulphate. The time required was less than 2 hours and the final yield 90%.

More recently, G. M. MizanurRahman and et al.(31) synthesized the isotopically enriched (CH₃²⁰¹Hg⁺) by methylation of mercuric oxide (²⁰¹HgO) with tetramethyltin (CH₃)₄Sn). The resulting reaction mixture was then stirred for one hour at 60 °C in water bath and isolation of CH₃²⁰¹Hg⁺ from the reaction mixture in a triple extraction steps with toluene (4+3+3 ml) after cooled and the extracts combination were then washed with deionised water three times (4+ 3 + 3 ml). Finally 2.5 ml of the toluene extract was dried as same as in R. C. Rodriguez Martin-Domimeadios. et al synthesis method (28). The syntheses time required was one hour and final yield > 90%.

Therefore, the main aim of this study was to produce of isotopically enriched monomethylmercury (²⁰¹MMHg) standard solution with minimum impurities, so as to achieve higher yield and shorter reaction time. For this purpose, the micro- scale preparation of isotopically enriched monomethylmercury (²⁰¹MMHg) procedure described in R. C. Rodriguez Martin-Domimeadios et al. (28) was applied to assess the possibility of producing micro-scale isotopically enriched methylmercury under standard laboratory conditions in order to use it as a standard compound for MMHg analysis in chlor alkali and gold mining hair samplers. The initial conditions were adapted from available literature methods according to the following criteria (28):

1. The isotopically enriched Mercury oxide (²⁰¹HgO) standard solution must be used as a starting material; this is the isotopically enriched from that is commercially available.

العدد الثاني عشر - ديسمبر 2016

2. The formation of dimethylmercury (DMHg) must be avoided.
3. Only small quantities of the starting material must be used, so it is necessary to prepare on a micro scale, because the starting material is highly cost.

The reaction product have been analysed and characterized by capillary gas chromatography (GC) coupled to inductively coupled plasma mass spectrometry (ICP-MS) for propylated species.

Experimental

Instrumentation

For speciated isotope dilution analysis (SIDMS) a gas chromatograph (GC) model HP 6850 equipped with a capillary column was coupled to an Agilent model HP-7500 ICP mass spectrometer via a heated steel transfer capillary as shown in **Figure 1**. The heated steel transfer capillary was inserted into the ICP torch injector and connection to the torch was realized by means of a glass T-piece. A conventional Meinhard concentric nebulizer and low volume water cooled cyclonic spray chamber were connected to the heated steel transfer capillary line connected ICP torch and this enabled continuous aspiration of a standard thallium solution ($25\mu\text{g l}^{-1}$). This configuration allowed optimization of instrument performance and simultaneous measurement of ^{203}Tl and ^{205}Tl for mass bias correction during the chromatographic run (18).




Figure 1 Schematic diagram for Agilent's GC-ICP-MS interface for 6850 GC/HP with 7500 ICP-MS/HP

The optimised operating conditions for the GC-ICP-MS coupling system are listed in Table 1

Table 1: GC and ICP-MS operating parameters.

العدد الثاني عشر - ديسمبر 2016

ICP-MS Instrument	Agilent 7500 series
Hg isotope acquired	196, 198, 199, 200, 201, 202, 204
Acquired mode	Time resolved
Dwell time	0.035 sec/point
RF power	1380 W
RF matching	1.53 V
Sample depth	7.2 mm
Torch-H	1.3 mm
Torch-V	0.5 mm
Carrier gas	Argon / 0.76 l min ⁻¹
Makeup gas	Argon / 0.22 l min ⁻¹
Extract 1	-2 V
Internal standard	Tl (25 ppb)
Nebulizer pump flow rate	0.20 rps
Spray chamber temperature	2 °C
GC Instrument	Agilent HP 6850
Injection	splitless – 1 µl
Oven program	50°C (1 min), 50°C/min →220°C (5 min)
Carrier gas	Helium
Transfer line temp	220°C
GC injector temp	220°C

Reagents and standards

All chemicals and used were of analytical reagent grade unless stated otherwise. Methanol, sodium acetate, acetic acid glacial (super grade) were purchased from VWR (BDH, UK). Sodium tetra-n-propylborate (NaBpR₄, ≥ 98% purity) was purchased from Chemos GmbH (Germany). Isooctane, Methylcobalamin and methylmercury (II) chloride were purchased from Sigma Aldrich (UK). Natural isotopic composition of inorganic mercury (In-Hg) standard stock solution for ICP (934 ±3.0 mg/kg) was purchased from Fluka (UK). Stock solution of methylmercury chloride (9543 mg/kg as Hg²⁺) of natural isotopic composition was prepared by dissolving methylmercury chloride (Sigma) in methanol (VWR). Working standard solutions were prepared fresh daily by appropriate dilution of the stock standard solutions in 1% HNO₃ and were stored in the fridge. Methylcobalamin (Sigma) used for synthesis was prepared by dissolution in an acetic acid- acetate buffer solution (0.1M, pH 5). Stock of isotopically enriched Mercury oxide (²⁰¹HgO) standard solution (994 mg/kg as ²⁰¹Hg) was prepared from mercuric oxide obtained from Oak Ridge National Laboratory

العدد الثاني عشر - ديسمبر 2016

(OaK, Ridge, USA). A part from toluene and isopropanol, which were HPLC grade were used. Double distilled water was used throughout.

Procedures:

Synthesis and primary analysis of ^{201}Hg – enriched methylmercury

A flow chart diagram of enriched monomethylmercury (MM^{201}Hg) synthesis protocol is shown in **Figure 1** below.




Figure 1 Flow chart for monomethylmercury MMHg isotopically enriched synthesis

Approximately 150 μl from 994 mg/kg ^{201}Hg –enriched inorganic mercury stock solution was transferred to the amber micro-reactors (Supelco) and diluted with 500 μl of buffer acetate (0.1M, pH 5). 3.09 mg of methylcobalamin was dissolved in another 3000 μl of buffer acetate (0.1M, pH 5) and 500 μl from this solution was added to the inorganic mercury solution. The pH was adjusted to pH 5 with concentrated sodium acetate solution. The solution was left sitting in the dark at 37 °C for 1 h. In order to stop the methylating reaction and to convert the unintentionally formed DMHg into methylmercury, the mixture was cooled at 4 °C, and 1.0 ml of concentrated hydrochloric acid was added and shaken for 5 minute. The MMHg formed was extracted three times with 500 μl of toluene. The combined toluene extracts were dried over sodium sulphate. 100 μl of this primary solution were diluted with 10 ml of isopropanol. Working solutions were prepared fresh daily by diluting the secondary isopropanol stock solution with deionized water as needed. The toluene extracts

العدد الثاني عشر - ديسمبر 2016

and the subsequent diluted solutions were stored in the fridge and protected from light until use.

Determination of the isotopic abundance and atomic weight concentration in the synthesised Me²⁰¹Hg enriched standard solution.

In order to calculate the purity of the synthesised enriched Me²⁰¹Hg standard solution and its atomic weight, 50 ppb of enriched Me²⁰¹Hg was prepared from stock solution (1.0 mg/kg enriched Me²⁰¹Hg secondary standard solution). The 1.0 ml of 50 ppb ($\mu\text{g kg}^{-1}$) was transfer to two clean glass vials and 5.0 ml of buffer acetate (0.1M, pH 3.9) was added in the glass vials and the pH adjusted at about 3.9. After which, 1.0 ml of iso-octane was added and propylated with 1.0 ml of 1% NaBPr₄. Then, Extraction of derivatised Hg species (peralkylated Hg) was done by vigorous shaking for 5 min and the iso-octane extract was afterwards centrifuged for 10 min at 3000 rpm then the extracted Hg species into iso-octane layer were transferred to GC vials and analysed with a coupling of GC-ICP-MS.

Reversed Isotope dilution mass spectrometry (RIDMS) method

To calculate the actual concentration of synthesised enriched Me²⁰¹Hg standard solution the RIDMS measurement was done by preparing of 1.0 mg kg⁻¹ enriched Me²⁰¹Hg and 1.0 0 mg kg⁻¹ normal abundance Me-Hg from synthesised enriched Me²⁰¹Hg and normal abundance MeHg stock solutions respectively. Then 100 μl from 1.0 0 mg kg⁻¹ enriched MeHg²⁰¹ was weighed in two clean and dry glass vials and spiked with 300 μl from 1.0 mg kg⁻¹ Me-Hg normal abundance and weighed again. 5.0 ml of 0.1 M buffer acetate (pH 3.9) was added in vials and the pH adjusted at about 3.9. Then, the derivatization and the Extraction of derivatised Hg species (peralkylated Hg) were done as same in section 2.3.2 above.

The concentration of the enriched methylmercury in the spiked solution and the reaction yield were calculated using reversed Isotope Dilution Mass Spectrometry (RIDMS) method by applying the isotope dilution equation (19).

$$c = \frac{C' W' A (RY' - X')}{w A' (X - RY)} \quad (1)$$

Where:

w = weight of the ²⁰¹Hg enriched solution (g)

c = Hg Concentration of the ²⁰¹Hg enriched solution (mg/kg)

A = Atomic mass (Hg) of the ²⁰¹Hg enriched solution (g/mole, from abundance det.)

Y = Isotope abundance of ²⁰¹Hg in the ²⁰¹Hg enriched solution (% , from abundance det.)

X = Isotope abundance of ²⁰²Hg in the ²⁰¹Hg enriched solution (% , from abundance det.)

W' = weight of the sample solution (g)

A' = Atomic mass (Hg) of the natural standard solution = 200.59 g/mole

Y' = Isotope abundance of ²⁰¹Hg in the natural standard solution = 13.2 %

X' = Isotope abundance of ²⁰²Hg in the natural standard solution = 29.8 %

العدد الثاني عشر - ديسمبر 2016

R = Isotope ratio of the integrated peak areas $^{202}\text{Hg}/^{201}\text{Hg}$ from IDMS measurements

C' = Hg concentration in the sample solution ($\mu\text{g}/\text{kg}$)

Therefore, the final parameter that requires measurement is the isotope ratio, $R(^{202}\text{Hg}/^{201}\text{Hg})$. A part from R , the other parameters is constants, masses and the concentration of the spike solution (normal abundance MeHg). Thus, the concentration (c) of the synthesised enriched Me ^{201}Hg standard solution is ultimately based on the measurement of isotope ratio (R) and the specification of the spiked material (normal abundance MeHg).

Results and discussion

Isotopically enriched MM ^{201}Hg synthesis and characterization

Isotopically enriched MMHg standard solution was synthesized from inorganic ^{201}Hg enriched stock standard solution according to the method described by R. C. Rodriguez Martin-Domimeadios et al (28). The analysis of the reaction products are then performed by CGC-ICP-MS after propylation with NaBPr $_4$. **Figure 3** shows the chromatogram obtained for mercury isotopes as can be seen; very low levels of inorganic mercury (Hg^{2+}) are present in the final product of MMHg. In addition, it can be observed very low levels of MeEtHg artifact from enriched MMHg during derivatization using NaBPr $_4$.




Figure 3 Chromatogram for mercury isotopes composition in synthesised enriched monomethylmercury ((MM ^{201}Hg)).

العدد الثاني عشر - ديسمبر 2016

The isotope composition of the enriched monomethylmercury (MM^{201}Hg) is shown in **Table 2**, and was calculated from measured peak area. This standard solution presents a $^{202}\text{Hg}/^{201}\text{Hg}$ ratio of 0.02510, which is highly different from the natural $^{202}\text{Hg}/^{201}\text{Hg}$ ratio of 2.2587. Therefore, the synthesised enriched MM^{201}Hg can be used directly as certified isotopically enriched Me^{201}Hg standard solution to perform isotope dilution (IDMS) analysis of MMHg in real samples (spiked solution).

Table 2: The atom fraction composition of enriched MM^{201}Hg aliquot, mass fractions and isotope ratio ($^{202}\text{Hg}/^{201}\text{Hg}$) determined by ICP-MS of the isotopically enriched methylmercury (Me^{201}Hg)

Hg isotope	Accurate mass	Abundance (mol at %)	
		Natural (IUPAC) ⁽³²⁾	*Enriched MM^{201}Hg
^{196}Hg	195.965	0.1534	0.000
^{198}Hg	197.967	9.9680	0.0604 ± 0.002
^{199}Hg	198.968	16.873	0.1295 ± 0.010
^{200}Hg	199.968	23.096	0.9071 ± 0.030
^{201}Hg	200.970	13.181	96.3856 ± 0.002
^{202}Hg	201.971	29.863	2.4172 ± 0.05
^{204}Hg	203.974	6.865	0.0994 ± 0.002
Sum (%)		100	100 ± 0.014
Hg atomic weight (g/mole)		200.598	201.014
R (202/201)		2.2587	0.02510

*Mean value \pm SD(n = 5)

Moreover, the actual concentrations of the enriched MM^{201}Hg in the spiked solution and in the reaction yield obtained were calculated using reverse isotope dilution (RIDMS) analysis (100 ul of enriched $1.0 \text{ mg kg}^{-1} \text{ MM}^{201}\text{Hg}$ standard solution spiked with 300 ul of 1.0 mg kg^{-1} normal abundance MeHg standard solution). Two independent IDMS experiments were carried out and the extracted Hg species from each solution was injected five times and the typical chromatogram obtained for mercury isotope showed in **Figure 4**. The average concentration of spiked solution turned out to be $1.31 \pm 0.04 \mu\text{g g}^{-1}$ and the methylation yield is 96.38 %. In addition, the stability of this enriched standard solution has been checked after 8 months and the degradation of about 25.23% was found.

العدد الثاني عشر - ديسمبر 2016




Figure 4 Chromatogram obtained from Reversed Isotope dilution mass spectrometry (RIDMS) experiment of synthesised ^{201}Hg – enriched monomethylmercury

Conclusion

A highly pure isotopically enriched monomethylmercury (MM^{201}Hg) has been synthesized from inorganic mercury (^{201}HgO) standard stock solution and methylcobalamin (MeCo) with a yield of more than 95% in a synthesis procedure lasting less than 1.5 hour at 37°C . The product is highly pure and stays stable for long time in cold conditions (-20°C). However, if this enriched solution is going to be used as standard, the concentration should be checked frequently. This isotopically enriched monomethylmercury (MMHg) standard solution has been successfully used for isotope dilution analysis of methylmercury in biological (hair) samples from contaminated area (chlor alkali plant and gold mining areas), environmental and food samples.

References

- (1) Li P, Feng X, Qiu G, Shang L, Li G. Human hair mercury levels in the Wanshan mercury mining area, Guizhou Province, China. *Environ Geochem Health* 2009;31 (6): 683-691.
- (2) Zhang Z, Wang Q, Zheng D, Zheng N, Lu X. Mercury distribution and bioaccumulation up the soil-plant-grasshopper-spider food chain in Huludao City, China. *Journal of Environmental Sciences* 2010 8;22 (8): 1179-1183.
- (3) Rahman GMM, Fahrenholz T, Kingston HM. Application of speciated isotope dilution mass spectrometry to evaluate methods for efficiencies, recoveries, and quantification of mercury species transformations in human hair. *J Anal At Spectrom* 2009;24 (1): 83-92.
- (4) Delgado A, Prieto A, Zuloaga O, de Diego A, Madariaga JM. Production of artifact methylmercury during the analysis of certified reference sediments: Use of ionic exchange in the sample treatment step to minimise the problem. *Anal Chim Acta* 2007 1/16;582 (1): 109-115.
- (5) Morita M, Yoshinaga J, Edmonds JS. The determination of mercury species in environmental and biological samples (Technical report). *Pure and Applied Chemistry* 1998;70 (8): 1585-1615.
- (6) Leermakers M, Baeyens W, Quevauviller P, Horvat M. Mercury in environmental samples: Speciation, artifacts and validation. *TrAC - Trends in Analytical Chemistry* 2005;24 (5): 383-393.
- (7) Kim SA, Jeon CK, Paek DM. Hair mercury concentrations of children and mothers in Korea: Implication for exposure and evaluation. *Sci Total Environ* 2008 8/25;402 (1): 36-42.
- (8) Agusa T, Kunito T, Iwata H, Monirith I, Tana TS, Subramanian A, et al. Mercury contamination in human hair and fish from Cambodia: levels, specific accumulation and risk assessment. *Environmental Pollution* 2005 3;134 (1): 79-86.
- (9) Chen S, Chou S, Hwang D. Determination of methylmercury in fish using focused microwave digestion following by Cu²⁺ addition, sodium tetrapropylborate derivatization, n-heptane extraction, and gas chromatography–mass spectrometry. *Journal of Chromatography A* 2004 1/23;1024 (1–2): 209-215.
- (10) Mizanur Rahman GM, Skip Kingston HM, Bhandari S. Synthesis and characterization of isotopically enriched methylmercury (CH₃201Hg⁺). *Applied Organometallic Chemistry* 2003;17 (12): 913-920.
- (11) Clarkson TW, Magos L. The toxicology of mercury and its chemical compounds. *Crit Rev Toxicol* 2006;36 (8): 609-662.
- (12) Li P, Feng X, Qiu G, Shang L, Wang S. Mercury exposure in the population from Wuchuan mercury mining area, Guizhou, China. *Sci Total Environ* 2008 6/1;395 (2–3): 72-79.
- (13) Holmes P, James KAF, Levy LS. Is low-level environmental mercury exposure of concern to human health? *Sci Total Environ* 2009;408 (2): 171-182.
- (14) Thomas W. Clarkson, Laszlo Magos, and Gary J. Myers. Human Exposure to Mercury: The Three Modern Dilemmas. 2003;16:321-322, 324-343.

العدد الثاني عشر - ديسمبر 2016

- (15) United Nations U. Technical guidelines on the environmentally sound management of mercury wastes. 2010 3 March;fifth version:8-9,10-84.
- (16) Veiga MM, Nunes D, Klein B, Shandro JA, Velasquez PC, Sousa RN. Mill leaching: a viable substitute for mercury amalgamation in the artisanal gold mining sector? J Clean Prod 2009;17 (15): 1373-1381.
- (17) Bancon-Montigny C, Yang L, Sturgeon RE, Colombini V, Mester Z. High-yield synthesis of milligram amounts of isotopically enriched methylmercury (CH₃¹⁹⁸HgCl). Applied Organometallic Chemistry 2004;18 (2): 57-64.
- (18) Krupp EM, Donard OFX. Isotope ratios on transient signals with GC-MC-ICP-MS. Int J Mass Spectrom 2005;242 (2-3): 233-242.
- (19) Martín-Doimeadios RCR, Krupp E, Amouroux D, Donard OFX. Application of isotopically labeled methylmercury for isotope dilution analysis of biological samples using gas chromatography/ICPMS. Anal Chem 2002;74 (11): 2505-2512.
- (20) Tseng CM, Amouroux D, Brindle ID, Donard OFX. Field cryofocussing hydride generation applied to the simultaneous multi-elemental determination of alkyl-metal(loid) species in natural waters using ICP-MS detection. J Environ Monit 2000;2 (6): 603-612.
- (21) Gao Y, Shi Z, Long Z, Wu P, Zheng C, Hou X. Determination and speciation of mercury in environmental and biological samples by analytical atomic spectrometry. Microchemical Journal 2012 7;103 (0): 1-14.
- (22) Rodrigues JL, Alvarez CR, Fariñas NR, Berzas Nevado JJ, Barbosa Jr. F, Rodriguez Martín-Doimeadios RC. Mercury speciation in whole blood by gas chromatography coupled to ICP-MS with a fast microwave-assisted sample preparation procedure. J Anal At Spectrom 2011;26 (2): 436-442.
- (23) Clough R, Truscatt J, Belt ST, Hywel Evans E, Fairman B, Catterick T. Isotope Dilution ICP-MS for Speciation Studies. Applied Spectroscopy Reviews 2003 01/04; 2012/09;38 (1): 101-132.
- (24) Hill SJ, Pitts LJ, Fisher AS. High-performance liquid chromatography–isotope dilution inductively coupled plasma mass spectrometry for speciation studies: an overview. TrAC Trends in Analytical Chemistry 2000 0;19 (2–3): 120-126.
- (25) Yang L, Colombini V, Maxwell P, Mester Z, Sturgeon RE. Application of isotope dilution to the determination of methylmercury in fish tissue by solid-phase microextraction gas chromatography–mass spectrometry. Journal of Chromatography A 2003 9/5;1011 (1–2): 135-142.
- (26) Liang L, Lazoff S. Evaluation of the procedure for alkaline digestion solvent estimation for methyl mercury artifact formation. Talanta 1999;48 (1): 231-233.
- (27) Hintelmann H, Evans RD. Application of stable isotopes in environmental tracer studies - Measurement of monomethylmercury (CH₃Hg⁺) by isotope dilution ICP-MS and detection of species transformation. Fresenius J Anal Chem 1997;358 (3): 378-385.
- (28) Rodríguez Martín-Doimeadios RC, Stoichev T, Krupp E, Amouroux D, Holeman M, Donard OFX. Micro-scale preparation and characterization of isotopically enriched monomethylmercury. Appl Organomet Chem 2002;16 (10): 610-615.



العدد الثاني عشر - ديسمبر 2016

- (29) Rodríguez-González P, Marchante-Gayón JM, García Alonso JI, Sanz-Medel A. Isotope dilution analysis for elemental speciation: a tutorial review. *Spectrochimica Acta Part B: Atomic Spectroscopy* 2005 2/28;60 (2): 151-207.
- (30) Rouleau C, Block M. Fast and high-yield synthesis of radioactive $CH_3^{203}Hg(II)$. *Applied Organometallic Chemistry* 1997;11 (9): 751-753.
- (31) Rahman GMM, Kingston HMS, Bhandari S. Synthesis and characterization of isotopically enriched methylmercury ($CH_3^{201}Hg^+$). *Appl Organomet Chem* 2003;17 (12): 913-920.
- (32) Michael B, Michael E. Wieser. Isotopic compositions of the elements 2009 (IUPAC Technical Report). 2011;83 (2): 397-410.