



## Libyan Journal of Public Health Practices (LJPHP)

Journal homepage:  
<https://journals.uob.edu.ly/LJPHP/index>



### Changes in Lifestyle and Eating Habits of the Libyan Population During the COVID-19 Pandemic

Souad F. El-Mani<sup>1</sup>, Ashmisa Eltuhami<sup>1</sup>, Reima M. Mansour<sup>1</sup>, Laila Elabbar<sup>2</sup>, Noha Bosrewill<sup>2</sup>, Shams Burwag<sup>2</sup>, Malak Alfadil<sup>2</sup>, Wafa Elabbar<sup>2</sup>

<sup>1</sup> Department of Nutrition, Faculty of Public Health, University of Benghazi, Benghazi, Libya

<sup>2</sup> Department of Nutrition, Faculty of Public Health, University of Benghazi, Benghazi, Libya

#### ARTICLE INFO

##### **Article history:**

Received 1/6/2024

Revised 10/6/2024

Accepted 24/6/2024

##### **Keywords:**

COVID-19

Eating habits

Physical activity

Meal consumption

Life style changes

#### ABSTRACT

The corona virus pandemic has had a massive impact on people globally. Due to national lockdowns and social distancing people all over the world have developed irregular lifestyles. The current study aimed to assess the lifestyle and eating habit changes of population during the COVID-19 pandemic. A cross-sectional study was conducted by random sampling (N =940) on the general population of the Libyan population during the period from February 9th, 2021 until March 15th, 2021. A questionnaire was used to assess the changes of lifestyle and eating habits during COVID-19, which later identified by analysis using the SPSS program. The majority of the respondents 81% were not infected with the corona virus, and less than half of them, 39% followed all the recommended procedures. To lower the risk of being infected there was an increase in supplement use among female participants in comparison to male participants (85% of female vs. 15%, male p.value =0.000). This study also indicated that most of participants had a sedentary life style and changes in eating habits. These changes were seen more among participants infected with corona virus than others who were not infected (51% vs. 45%, p=0.000). The current study concluded that there were changes in participants' life style during the COVID-19 pandemic; the majority of the participants had a sedentary lifestyle, and increase in their body weight and in meal consumption- especially during the evening Education programs should be conducted to promote healthy lifestyle and healthy eating to combat infections such as COVID-19.

## 1. Introduction

The coronavirus disease of 2019 is referred to as COVID-19, or Corona Virus. (1) The corona virus was first identified in Wuhan, China in December 2019, and due to its high contagiousness, it rapidly spread around the globe. (2) On January 31, 2020, the World Health Organization (WHO) designated COVID-19 as a public health emergency of global concern. A day later, on March 11, 2020, the WHO classified this epidemiological occurrence as a pandemic. (3,4) The WHO released a situation report on July 29, 2022, stating that there have been more over 579 million confirmed cases worldwide, with roughly 505,450 of those cases occurring in Libya. (5)

Although these limitations were implemented to decrease the spread of the COVID-19 virus, they also had a negative impact on mental and physical health in many nations and were found to have changed people's behavior, especially with regard to food habits. (10–13) Many studies have shown that when in stressful situations people tend to binge- resorting to unhealthy foods also known as “comfort food” - which are highly processed, sugar filled items that can have a deteriorating effect on health in the long term. (14, 15) In addition, due to the closure of gyms and the implementation of curfew, most people were forced to decrease their physical activity and have sedentary life style. (13–15) The impaired nutritional habits combined with decline levels of physical activity- could lead to a positive energy balance that may lead to weight gain and obesity. (16–19) Number previous studies reported that participants saw an increase in their meal number and frequency during lockdown. (20–22) In a study based in the Middle East and North Africa a total of 2970 participants were questioned. The percentage of participants consuming five or more meals per day increased from (2.2%) before the pandemic to (6.2%) during the pandemic. Moreover, the percentage of participants skipping meals decreased from (64.4%) before the pandemic to (45.1%) during the pandemic. When asked why they skipped

meals, (60.8%) stated that not having enough time was the main reason before the pandemic and (37.9%) stated that loss of appetite was the main reason for skipping meals during the pandemic. Although the results showed an increase in meal frequency, (74.0%) of participants reported not meeting the recommended water intake; drinking less than eight cups of water per day during the pandemic (12).

Due to the effect of the restriction on eating habits and behavior, the World Health Organization (WHO) released a list of nutritional recommendations that may enhance human immunity and decrease the complication of the disease. These recommendation suggested people should be follow a healthy diet that includes a variety of foods from each food group to ensure they are getting enough vitamins, minerals, dietary fiber, and protein, it also suggested people should limit sugar, fat, and salt intake. (23–29) Although these may not prevent people from getting COVID, they do help build a stronger immune system and lower the risk of chronic illness and severe complications.

There are limited studies that investigate the eating habits of people during the COVID-19 pandemic in Libya, and no study that has been undertaken in Benghazi, Libya, which explores the impact of COVID-19 on people's lifestyle and dietary patterns. Therefore, it has been suggested that the increase in home confinement and the psychological effect of the pandemic have induced changes in the lifestyle and eating patterns of individuals in Libya. The main aims of this study were to investigate the lifestyle and eating habit changes of participants during the COVID-19 pandemic and to determine the demographic variation in eating habits and lifestyle among the participants in Libya.

## 2. Methodology

**2.1. Study population:** A cross-sectional study was conducted by random sampling (N =940) on the general population of the Libyan population during the period from February 9<sup>th</sup>, 2021 until March 15<sup>th</sup>, 2021. The respondents were recruited to take part in this study either by

filling a questionnaire using Google form via online platform, or by the distribution of hard copies of the questionnaires to people in public spaces that include malls and schools with restricted by the safety procedures. The total number of participants who agreed to complete the questionnaire was N=940 (N=850 completed electronic questionnaire, and N=90 completed the hard copy).

**2.2. Questionnaire:** The study's questionnaire consisted of 38 questions separated into five sections. The first section covered demographics and personal information about the participants, such as city of residence, age, gender, educational level, occupation, income, and the number of family members. The second section of the questionnaire inquired about the medical history of the participants. The third section asked question about the changes in people's lifestyles during the COVID-19 pandemic. The fourth section asked questions regarding the changes in eating habits that occurred during COVID-19. The last section contained food frequency questionnaire. To reach the largest number of participants the questionnaire was distributed by two methods, in person and online through social media platforms such as twitter and facebook.

**2.3. Statistical Analysis:** The data was analyzed using SPSS version 21.0, the Statistical Package for the Social Sciences. In order to compare data, descriptive statistical tests such as frequencies, correlation tests, and Chi square tests with 95% confidence intervals were performed. Chi square analysis was employed to examine the relationship between different sociodemographic variables and the percentage of individuals who took supplements and/or herbs on a daily basis. In addition, the association between changes of life style, eating habits among participants and socio-demographic factors were tested using Chi square analysis. Correlation test was used to compare usual weight and current weight of participants. All p values <0.05 were considered statistically significant.

**2.4. Ethical Statement:** This study was approved by the University of Benghazi. All surveys were anonymous and unidentifiable to

maintain the confidentiality of the data collected.

### 3. Results

#### 3.1 Demographical characteristics of participants:

The questionnaire was completed by 940 participants; the majority of them 676(72%) were from Benghazi and (28%) from outside Benghazi including (Tripoli n=173, Misruta n=19, Albayda n=15, Almarj n=6, Sirt n=7, Kufrah n=11, Tobruk n=10, Zawra n=9, Hon n=10, Ajdabia =4 ). The majority of the surveyed individuals were female (78%) and the most distributed age was 20-35 (62%). Around half of the participants were single (56%), and the majority of them (85%) had university degree or higher level of education. There was a variation in jobs of the participants, (28%), (17%), and (12%) of them were students, teacher, and doctor respectively, while (10%) of participants were housewife and (6%) did not have any job. More than half of the respondents stated that their income was very good, and they had more than five members in their family, (58%) and (57%) respectively (**Table1**).

**Table 2** shows that the majority of participants (81%) were not infected by corona, while (19%) of them were infected by the corona virus. More than half of the participants (66%) had a family member infected by the corona virus. Most of the surveyed individuals (78%) reported following safety measurements during the COVID-19 pandemic. The majority of the respondents did not have any other disease and only small percentage of them reported having hearth problems those include (7%) hypertension (2%) heart disease (5%) diabetes and (8%) respiratory diseases. The majority of respondents (61%) reported using supplements during the pandemic, and half of the respondents (50%) reported using herbs during the pandemic. Female and participants of high level of education consumed more supplements than males and participants with low education levels (15% males vs. 85% females,  $p=0.000$ ; 12% primary and secondary level vs. 88% university and above level, respectively)

**Table 1: The Distribution of Demographical Characteristics of Participants**

| Characteristics                 | Frequency | Percentage |
|---------------------------------|-----------|------------|
| <b>Gender</b>                   |           |            |
| Female                          | 738       | <b>78%</b> |
| Male                            | 202       | 22%        |
| <b>Age</b>                      |           |            |
| <20                             | 73        | 8%         |
| 20-35                           | 584       | <b>62%</b> |
| 36-50                           | 204       | 22%        |
| 51-65                           | 63        | 7%         |
| >65                             | 12        | 1%         |
| <b>Location</b>                 |           |            |
| Benghazi                        | 680       | <b>72%</b> |
| Outside of Benghazi             | 260       | 28%        |
| <b>Education level</b>          |           |            |
| Primary level                   | 23        | 3%         |
|                                 | 113       | 12%        |
| Secondary level                 | 795       | <b>85%</b> |
| University and above            |           |            |
| <b>Social Status</b>            |           |            |
| Single                          | 529       | <b>56%</b> |
| Married                         | 370       | 40%        |
| Divorced                        | 24        | 3%         |
| Widowed                         | 12        | 1%         |
| <b>Occupation</b>               |           |            |
| Student                         | 267       | <b>28%</b> |
| Teacher                         | 169       | <b>17%</b> |
| Doctor                          | 108       | 12%        |
| Engineer                        | 65        | 7%         |
| Housewife                       | 90        | 10%        |
| Government                      | 78        | 8%         |
| Employed                        | 117       | 12%        |
| Other                           | 55        | 6%         |
| Don't have                      |           |            |
| <b>Income</b>                   |           |            |
| Good                            | 240       | 27%        |
| Very good                       | 519       | <b>58%</b> |
| Excellent                       | 133       | 15%        |
| <b>Number of Family Members</b> |           |            |
| ≤5                              | 396       | 43%        |
| >5                              | 518       | 57%        |

**3.2 Medical history of participants****Table 2: Medical history of participants:**

| Medical History                     | N   | %          |
|-------------------------------------|-----|------------|
| <b>Corona infection:</b>            |     |            |
| Yes                                 | 197 | 19%        |
| No                                  | 753 | <b>81%</b> |
| <b>Family corona infection:</b>     |     |            |
| Yes                                 | 317 | <b>66%</b> |
| No                                  | 611 | 34%        |
| <b>Following safety procedures:</b> |     |            |
| Yes                                 | 727 | <b>78%</b> |
| No                                  | 205 | 22%        |
| <b>Hypertension</b>                 |     |            |
| Yes                                 | 65  | 7%         |
| No                                  | 814 | <b>93%</b> |
| <b>Heart disease</b>                |     |            |
| Yes                                 | 17  | 2%         |
| No                                  | 847 | <b>98%</b> |
| <b>Diabetes</b>                     |     |            |
| Yes                                 | 43  | 5%         |
| No                                  | 829 | <b>95%</b> |
| <b>Respiratory disease</b>          |     |            |
| Yes                                 | 69  | 8%         |
| No                                  | 809 | <b>92%</b> |
| <b>Supplements intake:</b>          |     |            |
| Yes                                 | 552 | <b>61%</b> |
| No                                  | 359 | 39%        |
| <b>Herbs use:</b>                   |     |            |
| Yes                                 | 454 | <b>50%</b> |
| No                                  | 454 | 50%        |

**3.3 Participants' Life style changes during corona:**

When are asked the participants to describe the changes in their lifestyles during corona, approximately half of respondents 49% reported that they did not change in lifestyle and only 36% of them followed healthy life style. Nearly half of participants (43%) also state that they had changes in their health status and (58%) of them did not change the exercise level and only 16% had an increase exercise. Nearly half of the participants (47%) reported an increase in sleeping hours, and the majority sleeping hours were from 6-8 hours daily (49%). The same percentage of the surveyed individuals (54%) spent more than four hours on the television, and

they reported an increase in their weight during the pandemic (54%). In table 3 there was significant relation between the changes of sleeping hours and participants who are infected with corona. Patients with Covid 19 had changes in their health status and increase sleeping hours. (57% vs 40%,  $p=0.000$  and 61% vs 43%  $p=0.000$ ) than uninfected participants.

**Table 3: participants' life style changes during the pandemic of corona**

|                                 | N   | %   |
|---------------------------------|-----|-----|
| <b>life style changes:</b>      |     |     |
| Yes, became healthy             | 325 | 36% |
| Yes, became unhealthy           | 131 | 15% |
| Did not change                  | 443 | 49% |
| <b>Health status changes:</b>   |     |     |
| Yes                             | 388 | 43% |
| No                              | 508 | 57% |
| <b>Exercise changes:</b>        |     |     |
| Yes, increase exercise          | 128 | 16% |
| Yes, decrease exercise          | 210 | 26% |
| Did not change                  | 477 | 58% |
| <b>sleeping hours changes:</b>  |     |     |
| Yes, increase in sleeping hours | 425 | 47% |
| Yes, decrease in sleeping hours | 131 | 14% |
| Did not change                  | 357 | 39% |
| <b>Hours of sleeping</b>        |     |     |
| Less than 6 hours               | 119 | 13% |
| 6-8 Hours                       | 442 | 49% |
| More than 8 Hours               | 348 | 38% |
| <b>Hours of watching TV:</b>    |     |     |
| 0-2                             | 173 | 19% |
| 2-4                             | 243 | 27% |
| >4                              | 494 | 54% |
| <b>Weight changes</b>           |     |     |
| Increase in weight              | 493 | 54% |
| Decrease in weight              | 134 | 15% |
| Weight stayed the same          | 284 | 31% |

### 3.4 Food habits and food consumption among participants during the pandemic of corona virus:

Table 4 shows that around half of participants (46%) reported an increase in meal

consumptions, 62% of them consumed less than three meals per day and 40 % of them consumed two snacks per day. The most meal neglected among participants was breakfast (44%), and the most type of snack consumption were chocolate and pastries, 50% and 24%. More than half of respondents state that their meal consumption was increased at night and in the evening, 52% 54%, respectively.

**Table 4: Food habits and food consumption among participants during corona pandemic**

|                                    | N   | %   |
|------------------------------------|-----|-----|
| <b>Meal consumption changes:</b>   | 423 | 46% |
| Increase                           | 135 | 15% |
| Decrease                           | 358 | 39% |
| Did not change                     |     |     |
| <b>Number of meal consumption</b>  |     |     |
| 1-3                                | 569 | 62% |
| 3-6                                | 334 | 36% |
| More than 6                        | 16  | 2%  |
| <b>Number of snack consumption</b> |     |     |
| One snack                          | 310 | 34% |
| Two snacks                         | 363 | 40% |
| Three snacks                       | 166 | 18% |
| More than three snacks             | 69  | 8%  |
| <b>Type of snack consumption</b>   |     |     |
| Sweets                             | 78  | 9%  |
| Chocolate                          | 424 | 50% |
| Pasta                              | 204 | 24% |
| Nuts                               | 97  | 12% |
| Others                             | 43  | 5%  |
| <b>Neglected meals</b>             |     |     |
| Breakfast                          | 401 | 44% |
| Lunch                              | 72  | 8%  |
| Dinner                             | 188 | 21% |
| Snacks                             | 176 | 19% |
| Did not skip any meal              | 75  | 8%  |
| <b>Consumption increased:</b>      |     |     |
| <b>In the morning</b>              |     |     |
| Yes                                | 116 | 17% |
| No                                 | 551 | 83% |
| <b>In the afternoon</b>            |     |     |
| Yes                                | 171 | 26% |
| No                                 | 492 | 74% |
| <b>In the evening</b>              |     |     |
| Yes                                | 410 | 54% |
| No                                 | 350 | 46% |
| <b>At night</b>                    |     |     |
| Yes                                | 396 | 52% |
| No                                 | 364 | 48% |

During the pandemic, there was variation in types of food consumption. Approximately near to half of participant had an increase consumption of fruit and vegetable, tea and coffee and starchy food, 46%, 42% and 32% respectively. Whereas 30% of the participants had decrease fast food consumption and 67%, 70% of them reported that they did not change meat and restaurants food consumption **Table 5**.

#### 4. Discussion:

This study indicates that there were no changes in participants' lifestyle during the pandemic of COVID-19. That means the level of physical activity was already low before the pandemic,

therefore the pandemic had no effect on their sedentary life style. The present study's results are inconsistent with the findings from previous studies, which found that working from home, limitations on exercising, and gym closures during the COVID-19 pandemic were the main causes of the decline in physical activity levels (12, 30- 32).

The findings of this study revealed that more than half of the participants spent more than four hours watching television, similar to previous studies (31-36) which reported an increase of screen times in all age groups. This might be because participants were spending more time at home or the feeling of emptiness

**Table 5- Changes in type of food consumption during COVID-19**

|                                                   | N   | %  |
|---------------------------------------------------|-----|----|
| <b>Change in Meat Consumption</b>                 |     |    |
| Increase consumption                              | 145 | 17 |
| Decrease consumption                              | 144 | 16 |
| Did not change consumption                        | 587 | 67 |
| <b>Change in Fruit and Vegetable consumption</b>  |     |    |
| Increase in consumption                           | 406 | 46 |
| Decrease in consumption                           | 88  | 10 |
| No change in consumption                          | 357 | 44 |
| <b>Change in Tea and Coffee Consumption</b>       |     |    |
| Increase in consumption                           | 373 | 42 |
| Decrease in consumption                           | 119 | 14 |
| No change in consumption                          | 389 | 44 |
| <b>Changes in Fast Food Consumption</b>           |     |    |
| Increase in consumption                           | 246 | 29 |
| Decrease in consumption                           | 262 | 30 |
| No change in consumption                          | 357 | 41 |
| <b>Change in Starchy Food Consumption</b>         |     |    |
| Increase in consumption                           | 281 | 32 |
| Decrease in consumption                           | 154 | 18 |
| No change in consumption                          | 429 | 50 |
| <b>Increase in Consumption of Restaurant Food</b> |     |    |
| Increase in consumption                           | 267 | 30 |
| No change in consumption                          | 639 | 70 |

that occurred during the corona pandemic- which lead to people trying to find comfort through television. Other studies indicated that the sleep quality decreased and the participants were suffering from insomnia during the pandemic. (37,38) In the previous study nearly half of participants showed an increase in sleeping hours (6-8 hours daily).(37) The

findings of this study are similar with previous data obtained from China which confirmed total sleep time increased but sleep efficiency decreased. (34) Although the quality of sleep is not examined in this study, data from another Libyan study found that one-third of the Libyan population was suffering from clinical insomnia

during the pandemic, and there were changes in time and pattern of sleeping. (39)

**Table 5: The association between changes in sleeping hours , meal consumption and socio-demographicalf**

| <b>Variables</b>                                                                                | <b>Changes in sleeping hours</b>                      | <b>p.value</b> | <b>Changes meal consumption</b>                   | <b>p. value</b> |
|-------------------------------------------------------------------------------------------------|-------------------------------------------------------|----------------|---------------------------------------------------|-----------------|
| <b>Sex: #</b><br><b>Male</b><br><b>Female</b>                                                   | 77(18%)<br><b>346(82%)</b>                            | P=0.020#       | 71(17%)<br><b>350(83%)</b>                        | P=0.006#        |
| <b>Age: #</b><br><b>&lt;20</b><br><b>20-35</b><br><b>36-50</b><br><b>&gt; 50</b>                | 44(10%)<br><b>305(72%)</b><br>57(13%)<br>19(5%)       | P=0.000#       | 35 (8%)<br>298(71%)<br>71(17%)<br>18(4%)          | P=0.000#        |
| <b>Education level:</b><br><b>Primary and Secondary</b><br><b>University and above</b>          | 10 (2%)<br>414 (98%)                                  | P=0.98         | 9 (2%)<br>410 (98%)                               | P=0.52          |
| <b>Social status: #</b><br><b>Single</b><br><b>Married</b><br><b>Divorced</b><br><b>Widowed</b> | <b>289(68.4%)</b><br>129(30.4%)<br>3(0.7%)<br>2(0.5%) | P=0.000#       | <b>261(62%)</b><br>150(36%)<br>7(1.3%)<br>3(0.7%) | P=0.014#        |
| <b>Income:</b><br><b>Good</b><br><b>Very good</b><br><b>Excellent</b>                           | 111(28%)<br>230(58%)<br>57(14%)                       | P=0.94         | 113(28%)<br>229(57%)<br>57(14%)                   | P=0.913         |

# significant association p. value <0.05 by chi-square test

This study results found that an approximately half of the participants reported an increase in meal consumptions similar to previous studies which showed an increase in meal number and frequency. (18, 33). This could be due to sitting at home, feeling afraid and stressed which could lead participant to overeat. Data from different international countries concluded that home confinement due to COVID-19 pandemic lead to an adverse effect on mental well-being and emotional state. These effects were associated with unhealthy lifestyle behaviours including unhealthy eating habits (18).

Regarding eating habit, this study indicates that most of the participants had unhealthy eating habits which included an increase in consumption of meals in the evening and at night. These results were confirmed by earlier worldwide studies (18,30,40-42) that demonstrated an increase in meal consumption

and overeating among participants due to restrictions and stressful feelings. Although there is an increase in consumption of meals, this study found that most participants neglected breakfast. This could be due to disturbed quality of sleep and loss of appetite- this was confirmed by outcomes in a previous study (17). This study also found most

participants consumed unhealthy snacks such as chocolate and pasta. Similar results came from other studies that indicated psychological distress due to the pandemic of COVID-19 was associated with emotional eating and a higher consumption of unhealthy snacks that include beverages, sweets, high-sugar foods, and salty foods (12,43). In contrast, a Turkish study found a decrease in consumption of some foods such as chips, biscuits, chocolates, and frozen foods (44).

## 5. Conclusion:

The current study concluded that there were changes in participants' life style during the COVID-19 pandemic; the majority of the participants had a sedentary lifestyle. More than fifty percent of the participants did not participate in any physical activity, and reported an increase in sleeping hours. The major change among participants was an increase in body weight and an increase in meal consumption- especially during the evening. These changes were seen more among participants who were infected with the corona virus. In addition, there was an increase in consumption of supplementation and herbs; females consumed more supplements and herbs when compared to males.

## Recommendations

1. Further longitudinal studies should be conducted to address these changes.
2. Establishing awareness programs aimed to promote healthy eating and ways to boost immunity to combat infections such as COVID-19.
3. Education programs should be carried out to educate people on supplements intakes and the benefits of them reducing the severity of COVID-19.

## Acknowledgments:

We gratefully acknowledged all volunteers who participated in this study, as well as my colleagues for assistance in completing this study.

## Conflict of interest:

Authors have declared that no conflict of interest

## References:

- 1- Vergnaud S. What does COVID-19 stand for?. Cited at: <https://www.goodrx.com/blog/what-does-covid-19-mean-who-named-it/>. Retrieved on: 26th May. 2020
- 2- de Lemos JA, McGuire DK, Drazner MH. B-type natriuretic peptide in cardiovascular disease. *The Lancet*. 2003 Jul 26;362(9380):316-22.
- 3- World Health Organization. COVID 19 Public Health Emergency of International Concern

- (PHEIC). Global research and innovation forum: towards a research roadmap.2020.
- 4- Watson OJ, Barnsley G, Toor J, Hogan AB, Winskill P, Ghani AC. Global impact of the first year of COVID-19 vaccination: a mathematical modelling study. *The Lancet infectious diseases*. 2022 Sep 1;22(9):1293-302
  - 5- World Health Organization. Libya /statistical [Internet]. [cited 2021 Apr 19]. Available from: <https://www.who.int/countries/lby/>
  - 6- Certain Medical Conditions and Risk for Severe COVID-19 Illness | CDC [Internet]. [cited 2021 Apr 18]. Available from: <https://www.cdc.gov/coronavirus/2019-ncov/need-extra-precautions/people-with-medical-conditions.html>
  - 7- Muniyappa R, Gubbi S. COVID-19 pandemic, coronaviruses, and diabetes mellitus. *American Journal of Physiology-Endocrinology and Metabolism*. 2020 May 1;318(5):E736-41.
  - 8- Petrakis D, Margină D, Tsarouhas K, Tekos F, Stan M, Nikitovic D, Kouretas D, Spandidos DA, Tsatsakis A. Obesity-a risk factor for increased COVID-19 prevalence, severity and lethality. *Molecular medicine reports*. 2020 Jul 1;22(1):9-19.
  - 9- Mueller AL, McNamara MS, Sinclair DA. Why does COVID-19 disproportionately affect older people?. *Aging (albany NY)*. 2020 May 5;12(10):9959.
  - 10- Eating well during Coronavirus / COVID-19 [Internet]. [cited 2021 Apr 18]. Available from: <https://www.bda.uk.com/resource/eating-well-during-coronavirus-covid-19.html>
  - 11- Balanzá-Martínez V, Atienza-Carbonell B, Kapczinski F, De Boni RB. Lifestyle behaviours during the COVID-19-time to connect. *Acta Psychiatrica Scandinavica*. 2020 May;141(5):399.
  - 12- Ismail LC, Osaili TM, Mohamad MN, Al Marzouqi A, Jarrar AH, Zampelas A, Habib-Mourad C, Jamous DO, Ali HI, Al Sabbah H, Hasan H. Assessment of eating habits and lifestyle during the coronavirus 2019 pandemic in the Middle East and North Africa region: a cross-sectional study. *British Journal of Nutrition*. 2021 Sep;126(5):757-66.
  - 13- Di Renzo L, Gualtieri P, Cinelli G, Bigioni G, Soldati L, Attinà A, Bianco FF, Caparello G, Camodeca V, Carrano E, Ferraro S. Psychological aspects and eating habits during COVID-19 home confinement: results of EHLCCOVID-19 Italian online survey. *Nutrients*. 2020 Jul 19;12(7):2152.
  - 14- Razzoli M, Pearson C, Crow S, Bartolomucci A. Stress, overeating, and obesity: Insights from human studies and preclinical models. *Neuroscience & Biobehavioral Reviews*. 2017 May 1;76:154-62.

- 15- Munsch S, Meyer AH, Quartier V, Wilhelm FH. Binge eating in binge eating disorder: a breakdown of emotion regulatory process?. *Psychiatry research*. 2012 Feb 28;195(3):118-24.
- 16- Górnicka M, Drywień ME, Zielinska MA, Hamułka J. Dietary and lifestyle changes during COVID-19 and the subsequent lockdowns among Polish adults: a cross-sectional online survey PLifeCOVID-19 study. *Nutrients*. 2020 Aug 3;12(8):2324.
- 17- Husain W, Ashkanani F. Does COVID-19 change dietary habits and lifestyle behaviours in Kuwait: a community-based cross-sectional study. *Environmental health and preventive medicine*. 2020 Dec;25(1):1-3.
- 18- Ammar A, Brach M, Trabelsi K, Chtourou H, Boukhris O, Masmoudi L, Bouaziz B, Bentlage E, How D, Ahmed M, Müller P. Effects of COVID-19 home confinement on eating behaviour and physical activity: results of the ECLB-COVID19 international online survey. *Nutrients*. 2020 May 28;12(6):1583.
- 19- Hill JO, Wyatt HR, Peters JC. Energy balance and obesity. *Circulation*. 2012 Jul 3;126(1):126-32.
- 20- Di Renzo L, Gualtieri P, Pivari F, Soldati L, Attinà A, Cinelli G, Leggeri C, Caparello G, Barrea L, Scerbo F, Esposito E. Eating habits and lifestyle changes during COVID-19 lockdown: an Italian survey. *Journal of translational medicine*. 2020 Dec;18(1):1-5.
- 21- Sidor A, Rzymiski P. Dietary choices and habits during COVID-19 lockdown: experience from Poland. *Nutrients*. 2020 Jun 3;12(6):1657.
- 22- Phillipou A, Meyer D, Neill E, Tan EJ, Toh WL, Van Rheen TE, Rossell SL. Eating and exercise behaviors in eating disorders and the general population during the COVID-19 pandemic in Australia: Initial results from the COLLATE project. *International Journal of Eating Disorders*. 2020 Jul;53(7):1158-65.
- 23- Davanzo R, Moro G, Sandri F, Agosti M, Moretti C, Mosca F. Breastfeeding and coronavirusdisease-2019: Adinterim indications of the Italian Society of Neonatology endorsed by the Union of European Neonatal & Perinatal Societies. *Maternal & Child Nutrition*. 2020 Jul;16(3):e13010.
- 24- Carver PE, Phillips J. Novel coronavirus (COVID-19): what you need to know. *Workplace health & safety*. 2020 May;68(5):250-.
- 25- Dietitians of Canada - Advice for the general public about COVID-19 [Internet]. [cited 2021 Jun 10]. Available from: <https://www.dietitians.ca/News/2020/Advice-for-the-general-public-about-COVID-19>.
- 26- World Health Organization. coronavirus disease (COVID-19): Pregnancy and childbirth [Internet]. [cited 2021 Jun 10]. Available from: <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/question-and-answers-hub/q-a-detail/coronavirus-disease-covid-19-pregnancy-and-childbirth>
- 27- UNICEF. Easy, affordable and healthy eating tips during COVID-19 | UNICEF [Internet]. [cited 2021 Jun 10]. Available from: <https://www.unicef.org/coronavirus/easy-affordable-and-healthy-eating-tips-during-coronavirus-disease-covid-19-outbreak>.
- 28- CDC. Coronavirus Disease (COVID-19) and Breastfeeding | Breastfeeding | CDC [Internet]. [cited 2021 Jun 10]. Available from: <https://www.cdc.gov/breastfeeding/breastfeeding-special-circumstances/maternal-or-infant-illnesses/covid-19-and-breastfeeding.html>
- 29- de Faria Coelho-Ravagnani C, Corgosinho FC, Sanches FL, Prado CM, Laviano A, Mota JF. Dietary recommendations during the COVID-19 pandemic. *Nutrition reviews*. 2021 Apr;79(4):382-93.
- 30- Phillipou A, Meyer D, Neill E, Tan EJ, Toh WL, Van Rheen TE, Rossell SL. Eating and exercise behaviors in eating disorders and the general population during the COVID-19 pandemic in Australia: Initial results from the COLLATE project. *International Journal of Eating Disorders*. 2020 Jul;53(7):1158-65.
- 31- Matsungo TM, Chopera P. Effect of the COVID-19-induced lockdown on nutrition, health and lifestyle patterns among adults in Zimbabwe. *BMJ Nutrition, Prevention & Health*. 2020 Dec;3(2):205.
- 32- Deschassaux-Tanguy M, Druesne-Pecollo N, Esseddik Y, De Edelenyi FS, Allès B, Andreeva VA, Baudry J, Charreire H, Deschamps V, Egnell M, Fezeu LK. Diet and physical activity during the coronavirus disease 2019 (COVID-19) lockdown (March–May 2020): Results from the French NutriNet-Santé cohort study. *The American journal of clinical nutrition*. 2021 Apr;113(4):924-38.
- 33- Górnicka M, Drywień ME, Zielinska MA, Hamułka J. Dietary and lifestyle changes during COVID-19 and the subsequent lockdowns among Polish adults: a cross-sectional online survey PLifeCOVID-19 study. *Nutrients*. 2020 Aug 3;12(8):2324
- 34- Li Y, Qin Q, Sun Q, Sanford LD, Vgontzas AN, Tang X. Insomnia and psychological reactions during the COVID-19 outbreak in China. *Journal of Clinical Sleep Medicine*. 2020 Aug 15;16(8):1417-8.
- 35- Wang X, Li Y, Fan H. The associations between screen time-based sedentary behavior and depression: a systematic review and meta-

- analysis. BMC public health. 2019 Dec;19(1):1-9.
- 36- Carroll N, Sadowski A, Laila A, Hruska V, Nixon M, Ma DW, Haines J, Guelph Family Health Study. The impact of COVID-19 on health behavior, stress, financial and food security among middle to high income Canadian families with young children. *Nutrients*. 2020 Aug 7;12(8):2352.
  - 37- Casagrande M, Favieri F, Tambelli R, Forte G. The enemy who sealed the world: effects quarantine due to the COVID-19 on sleep quality, anxiety, and psychological distress in the Italian population. *Sleep medicine*. 2020 Nov 1;75:12-20.
  - 38- Cellini N, Canale N, Mioni G, Costa S. Changes in sleep pattern, sense of time and digital media use during COVID-19 lockdown in Italy. *Journal of sleep research*. 2020 Aug;29(4):e13074.
  - 39- Elhadi M, Alsoufi A, Msherghi A, Alshareea E, Ashini A, Nagib T, Abuzid N, Abodabos S, Alrifai H, Gresea E, Yahya W. Psychological health, sleep quality, behavior, and internet use among people during the COVID-19 pandemic: a cross-sectional study. *Frontiers in psychiatry*. 2021 Mar 31;12:632496.
  - 40- Moynihan AB, Tilburg WA, Igou ER, Wisman A, Donnelly AE, Mulcaire JB. Eaten up by boredom: consuming food to escape awareness of the bored self. *Frontiers in psychology*. 2015 Apr 1;6:369.
  - 41- Di Renzo L, Gualtieri P, Pivari F, Soldati L, Attinà A, Cinelli G, Leggeri C, Caparello G, Barrea L, Scerbo F, Esposito E. Eating habits and lifestyle changes during COVID-19 lockdown: an Italian survey. *Journal of translational medicine*. 2020 Dec;18(1):1-5.
  - 42- Sidor A, Rzymski P. Dietary choices and habits during COVID-19 lockdown: experience from Poland. *Nutrients*. 2020 Jun 3;12(6):1657.
  - 43- Bermanian M, Mæland S, Blomhoff R, Rabben ÅK, Arnesen EK, Skogen JC, Fadnes LT. Emotional eating in relation to worries and psychological distress amid the COVID-19 pandemic: a population-based survey on adults in Norway. *International journal of environmental research and public health*. 2021 Jan;18(1):130.
  - 44- Özlem A, Mehmet N. COVID-19 Pandemi Kilitlenmesi Sürecinde Yeme Alışkanlıklarındaki Değişiklikler. *ESTÜDAM Halk Sağlığı Dergisi*. 2020;5:188-96.