

Petrographical and Paleontological Characterizations of the Quaternary Calcarenes along the Coastline of NE-Libya

Saad K. El Ebaidi¹, Mohammed H. Al Riaydh¹, Ahmed M. Muftah^{*1} and Hamed O. Elwerfalli²

¹ University of Benghazi, Faculty of Science, Department of Earth Sciences, Benghazi-Libya.

² Arabian Gulf Oil Company, Exploration Department, Geological Laboratory, Benghazi-Libya.

Received: 3/3/2020; accepted: 9/6/2020.

المخلص

ترسبات الكالكارينيت (الرمال الجيري المتناسك) التابع للعصر الرباعي يمكن مشاهدتها بشكل متقطع غير مستمر على طول الشاطئ الليبي بمياه البحر المتوسط، وكذلك الشاطئ التونسي، والابيطالي، والمالطي والقيصري. يتواجد الكالكارينيت في اماكن محدودة بالجبل الاخضر مكونا صخور متوازية مع الساحل الليبي وعلى ارتفاعات متفاوتة تصل أحيانا لمئات الأمتار فوق مستوى سطح البحر. تهدف هذه الورقة إلى وصف وتصنيف رسوبيات الكالكارينيت الرملية الكلسية في الحقبة الرباعية بالعصر الهلوسيني والبليستوسيني المتواجدة بالجبل الأخضر وبناء على النسيج الترسبي. تم أخذ عينات صخرية من أربع مناطق شاطئية بالمنطقة الشرقية وهي: قاريونس، والحنية، والحمامة، وسوسة. وفي المجمل تتكون هذه الصخور من رسوبيات بحرية بأعماق مائية مختلفة. وتتواجد هذه الصخور على شكل أحزمة متعاقبة كبيرة الحجم أفقية وعمودية وعلى هيئة أحواض مقعرة أو حادة من صخور ناعمة وأحيانا صلبة نوعا ما وبلون أبيض إلى كريم، مع تواجد قوالب الجذور والقواقع البرية في اماكن محدودة. اما الفتات الحيوي تمثله المنخربات القاعية الصغيرة الحجم التابعة للخزفيات والروتاليدات وكذلك التلازنيات، والامستراكودات، وبقايا القناذف البحري والرخويات والطحالب الجيرية الحمراء. بينما بموقع قاريونس فالصخور تتميز بتواجد غزير لصداف الرخويات المتمثلة في الكارديوم مع تخلل للمارل والطين الغني برخويات الأويستر.

الكلمات المفتاحية: الكالكارينيت، الجبل الاخضر، النسيج الصخري، الساحل الليبي، الطبقة المتقاطعة.

Abstract

Quaternary calcarenites deposits can be traced sporadically along the Mediterranean coastlines including Libya, Tunisia, Italy, Malta and Cyprus. Quaternary calcarenites are cropping out at local places in Al Jabal al Akhdar forming the coastal rocks, few hundred meters above sea level. This paper aimed to describe and classify the Quaternary (Pleistocene and Holocene) calcarenites of Al Jabal al Akhdar based on the depositional texture and the manifested sedimentary structures. Four rocky beach localities are subjected to sampling for this study namely, Garyounis, Al Haniyah, Al Hammamah and Susah. They are mainly composed of marine deposited at more or less various water depths. The concerned calcarenites displayed an alternating band of soft and hard white to creamy, planar to festoon trough large scale cross bedded with local occurrences of root casts and land snails. The bioclasts are represented by small benthic foraminifers of miliolids, rotalids and textularids as well as ostracodes, echinoderm remains, mollusks and calcareous red algae. The calcarenite in the studied localities showed wide range of facies distribution on both lateral and vertical sense in terms of biofacies and lithofacies, the calcarenite grains in Susah region is bounded by calcite bridges filled fractured. However, the calcarenite in Garyounis area is characterized by the presence of highly molluscan shells concentration such as *Cardium*, with intercalated marls and clays enriched with *Ostrea*.

Keywords: Calcarenite, cross bedded, Quaternary, Susah, Garyounis, Al Jabal al Akhdar

1. INTRODUCTION

The Quaternary calcarenite rocks in Libya was intensively studied (1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14). In general, the calcarenites are soft, highly porous sedimentary rocks formed by cementation of calcareous grains.

The Quaternary calcarenite rocks are well distributed in many places along Cyrenaica shoreline and (Fig. 1). Four localities are subjected to sampling that are; Al Haniyah, Al Hammamah, Susah and Garyounis calcarenites (Fig. 1). They revealed that these selected measured sections are mainly composed of marine bioclasts that deposited at more or less at various water depths. The concerned calcarenites displayed an alternating bands of soft and hard light brown to creamy planar to festoon

*Correspondence:

Ahmed M. Muftah
Department of Earth Sciences
University of Benghazi
ahmed.alkowafi@uob.edu.ly

trough large scale cross bedded with local occurrences of root casts and land snails. The hard beds of the Quaternary calcarenites were widely used in the era of Greek civilization in construction and that's adding historical event with a significant cultural value to this type of rock ¹⁵.

2. SEDIMENTOLOGY AND STRATIGRAPHY

2.1 Garyounis Calcarenite Petrography

The best developed calcarenites outcrop in southwest of Wadi Al Qattarah, Benghazi plain. These calcarenites unconformably overlies the Middle Miocene Benghazi Formation. The studied section is about 2 m thick, it consists of three main units ¹⁴ (Figs.2; 3 and 4).

concentration made of *Cerastoderma edula*, calcareous red algae, bryozoans and foraminifera.

Unit 2; consists of marls and clays, yellow to gray in color, soft, friable, thickly bedded (0.75 m), this unit also characterized by shell concentration made of *Ostrea* sp., the washed residue of this unit is made of ostracodes, *Elphidium macellum* and *Ammonia* cf. *tepida*.

Unit 3; yellow to brown calcarenite, moderately hard to hard and very thick bedded (1m), this unit is also characterized by shell concentration made of *Cerastoderma edula* Linnaeus.




Figure 1: Location map of the studied localities, NE Libya




Figure 2: The stratigraphical columnar sections of the Quaternary calcarenite rocks; A- Sush, B- Al Hammamah, C- Al Haniyah and Garyounis area

Unit 1; consists of calcarenite, yellowish white in color, hard, thickly bedded (0.3 m), this unit also characterized by shell

No fluid movement allowed due to lack of porosity because of dolomitization and cementation. The paleontological collections

from the Calcarenites in this study area are illustrated in Plate 1 and 2 and were based on the collection of the authors and ¹⁴.




Figure 3: Typical section of the calcarenite with high concentration of Oyster in the middle part (clay & marl) at Garyounis area.




Figure 4: The lower calcarenite (A & B); A) highly porous with thick layer of isopachous cement; B) Red algae; and the upper part (C & D); C) large thick-shelled gastropod; D) highly porous with thin layer of isopachous cement ¹⁴.

2.2 Petrography of Al Haniyah Calcarenites

The base of the Al Haniyah calcarenite overlies unconformably a pitted erosion surface of hard, white limestone bioturbated sediments with land snails. Two types of cross bedding, the lower one is planar and the upper one is festoon cross bedding (trough x-bedding). It composes of elongate semi ellipsoidal plunging trough or scoop like structures or hummocky cross bedding with total thickness of 24 m (Figs. 2 and 5). Thin sections show that there are three types of cement; isopachous, meniscus and the pore filled is blocky or drusy (Fig. 6). Mimetically, euhedral to anhedral dolomite crystals were observed (2 to 8 % max.), it occurs both as fabric selective replacement of the carbonate mud matrix and as a cement. Quartz, gypsum and iron minerals are also observed. Some of small benthic foraminifers are retrieved from this section such as *Lobatula lobatula*, *Elphidium crispum*, *E. advenum*, *E. cf. jensenii*, *Ammonia beccarii*, *Triloculina plicata*, *T. sp.*, *Quinqueloculina cf. bosciana.*, *Q. vulgarism*, *Spiroloculina angulosa*, *S. dilatata*, *Adelosina sp.*, *Cibicides cf. reflunens*, *Cibicides cf. variabilis*, *Rosalina bradyi*, *R. cf. macropora* and *Planorbulina mediterraneensis*. This assemblage is associated with rare ostracodes and small sized *Cerithium sp.*, bryozoan fragments and echinoderm spines. However, polychaetes serpulids are encrusted on some beds with terrestrial *Helix sp.*



Figure 5: Hummocky and Swale cross beddings in Al Haniyah calcarenite




Figure 6: A; shows micrite envelopes developed in algal peloids; there are three cement generations; B) isopachous cement; C) meniscus cement; and D) drusy or blocky cement, (40 x; PPL).

2.3 Al Hammamah Calcarenites

Al Hammamah calcarenite is unconformably overlies the Eocene carbonate rocks, bioclastic, and sandy limestone, brown to creamy in color. It consists of two types of cross bedding; planar (Figs. 2 and 7) and festoon cross bedding (trough x-bedding), with total thickness of 20 m (Fig. 8) with land snails (*Helix*) and root casts or molds (Fig. 9). Thin sections show that there are three types of cement; isopachous, meniscus and pore-filling as blocky or drusy (Fig. 10). Euhedral to anhedral dolomite crystals were observed (8 to 10 % max.), it occurs both as fabric selective replacement of the carbonate mud matrix and as cement (blocky cement is totally replaced by dolomite crystals). Quartz, gypsum and iron minerals are also observed. Al Hammamah in its type section is fossiliferous, notably of benthic foraminifers include *Quinqueloculina spp.*, *Elphidium advenum*, *E. crispum*, *Rosalina bradyi*, *Ammonia beccarii* and *Cibicides cf. reflunens*. This assemblage is associated with common small sized *Cerithium* with echinoderm spines and some terrestrial *Helix sp.* Present as well in places.



Figure 7: Planar cross beddings in Al Hammamah road-cut



Figure 8: Trough cross stratification in Al Hammamah area.




Figure 9: A) Horizontal Rhizoconcretion with *Helix* sp.; B) Tangential Rhizoconcretion (calcareous root concretions); C) encrusted Serpulid worm tubes; at Al Hammamah calcarenite.




Figure 10: A, B, C & D Shows meniscus cement in; B) isopachous cement; C) micrite envelopes developed in algae and some in miliolids, (40 x; PPL).

2.4 Susah Calcarenite

Susah calcarenite overlies unconformably an erosional surface of Eocene Apollonia rocks, bioclastic, and sandy limestone, brown to cream in color with total thickness of 18 m (Fig. 11). It consists of rhizoconcretions are parallel to the cross beddings (Figs. 2 and 12). Three types of cross bedding, planar and festoon cross bedding (trough x-bedding); large dune of hummocks have inside gently curve lamination which are convex up in the hummocks and there are also concave up lamination called swales between the hummocks (Fig. 13). Thin sections show that there are two types of cement; isopachous and meniscus (Fig. 14), euhedral to anhedral dolomite crystals were observed (of about 10 %), it occurs both as fabric selective replacement of the carbonate mud matrix and as a cement. Quartz, gypsum, anhydrite and iron minerals are also observed.



Figure 11: A and B) Cross bedding in Susah calcarenites; C) Hard beds of the Quaternary calcarenite were widely used in city construction of Greek civilization.




Figure 12: A & a1) Rhizoconcretions (biogenic), and plant root casts or molds are parallel to the cross stratifications.




Figure 13: A) Hummocky and Swale cross stratifications; B) Rhizoconcretions (biogenic), and plant root casts or molds




Figure 14: A, B & C) Red algae fragments, slightly compacted with meniscus cement; D) Isopachous cement, (40 x; PPL).

These Quaternary calcarenites are Pleistocene in age (Early Calabrian – Late Tarantian) according to.¹⁶ using ($^{87}\text{Sr}/^{86}\text{Sr}$) radiometric dating of the crystalline calcites from different localities.

3. DEPOSITIONAL ENVIRONMENT

Due to high diversity of bioclasts such as foraminifera (Plate 1), pelecypods (Plate 2), gastropods, bryozoans, echinoids, and

algae, the studied calcarenites are deposited in-situ at shallow marine carbonate environments (Fig. 15).

Al Haniyah, Al Hammamah and Susah are all generated by strongly sea water storm wave in the lower sea floor of shore-face. Most the calcarenites in NE Libya are subjected to meteoric water diagenesis and dissolution occurs in the phreatic environment; isopachous cement is formed below the water table, pores are totally filled with water and cementation continued and formed the drusy mosaic or blocky cements. The meniscus cement can precipitated particularly in the upper part of beach, occurred in the vadose environment. In the lagoon only mollusks, algae and foraminifera are noted. No fluid movement allowed due to lack of porosity because of dolomitization and cementation.

4. CONCLUSION

The Al Haniyah, Al Hammamah, and Susah calcarenites overlies unconformably a pitted erosional surface of Eocene limestone, while the Garyounis calcarenite overlies unconformably an erosion surface of middle Miocene of Benghazi carbonate rocks. Calcarenites may have been deposited on shallow banks with temporal periods of strong storm wave, and subsequently uplifted, they alternatively suggested as an aeolian deposit. The geographical matching similarity in Libyan coastal Quaternary outcrops, suggesting that Quaternary Cyrenaican calcarenite outcrops at Al Haniyah, Al Hammamah, and Susah could be stratigraphically assigned to Gergaresh Formation, however, Garyounis calcarenite stratigraphically assigned to Agedabia Formation. Al Haniyah, Al Hammamah and Susah were deposited by storm wave in the lower sea floor of shore-face shallow marine shelf.

ACKNOWLEDGEMENT

Thanks are due to Eddawi Ali Elhatmi the supervisor of the Geological Laboratory and Mahmoud Saleem the Exploration Manager of Libyan Petroleum Institute for the assistance in thin sections preparation.




Figure 15: Depositional schematic setting of the studied Quaternary calcarenites



Plate 1: 1) *Elphidium macellum* 60-70X; 2) *Elphidium advenum* 60-70X; 3) *Ammonia* cf. *beccari* 60-70X; 4) *Ammonia* cf. *tepida*; 5-6) Ostracod G1- spp. 60-70X. based on biota (after ¹⁴)



Plate 2: 1-2) *Venus* sp.; 3-4) *Tellina* sp.; 5-8) *Cerastoderma edula*; and 9-12) *Ostrea* sp.

6. REFERENCES

1. Stefanini G. Struttura geologica della Cirenaica e Cenn Descrittivi a corredo dello schizzo geologico dimostrativo della Cirenaica. In: La Cirenaica geografica, economics politica. Milano, 1923; 215-236.
2. Castany G. Les Extrusions Jurassique en Tunisie, Tunisia Serv. Mines, Ann. Mines et Geol., no. 14; 1955; 71 pp. illustr. Tunis.
3. Hey RW. The Quaternary and Paleolithic of northern Libya. *Quaternaria* 6; 1962: 435- 449.
4. Rohlich P. Geological Map of Libya, Explanatory Booklet Sheet: Al Bayda, NI 34-15, Libyan Arab Republic, Industrial Research Center. 1974
5. Klen L. Geological Map of Libya, 1:250,000. Sheet Benghazi XI 34-14, Explanatory' Booklet, Industrial Research Center, Tarabulus. 1974; 56 pp.,
6. Mann K. Geological map of Libya. 1:250,000 Sheet: Al Khums NI33-14 Explanatory Booklet. Industrial Research Centre, Tripoli, 1975a; 88 pp.
7. Mann K. Geological map of Libya. 1:250,000 Sheet: Misratah NI33-15 Explanatory Booklet. Industrial Research Centre, Tripoli, 1975b; 30 pp.
8. El Hinnawy M. and Cheshitev G. Geological map of Libya. 1:250,000 Sheet: Tarabulus NI33-13 Explanatory Booklet. Industrial Research Centre, Tripoli, 1975; 66 pp.
9. Mijalkovic N. Geological map of Libya, Explanatory Booklet Sheet: AL Qaddahiyah, NH 33-3, Libyan Arab Republic, Industrial Research Center. 1977.
10. Innocenti F. and Pertusati P. Geological map of Libya, Explanatory Booklet Sheet: AL Aqaylah, NH 34-5, Libyan Arab Republic, Industrial Research Center. 1984.
11. Al-Jamaily YW. and Abu-Arabyia, H. Microfossils study and paleoenvironmental conclusions of Gargaresh Formation (Quaternary), from selected exposed section, western beach of Zliten area (in Arabic), *Applied science*, 2000; 3: 5-12.
12. Minas HA. Palaeoenvironmental Reconstruction of The Gargaresh Formation, NW Libya, The geology of northwest Libya (sedimentary basins of Libya), 2003; Vol. II: 39-49.
13. Minas HA. and El Bakush HS. Shoreline Outcrops Comparison of Gargaresh Formation with Quaternary Calcarene Rocks of Cyrenaica, Northern Libya. Proceedings of the 5th International Conference on the Geology of Africa. 2007 (Oct. 2007); Vol. (1); P-P. Vi-71 – Vi-85 Assiut-Egypt.
14. Al Riaydh MH. and Abdulsamad EO. Agedabia Formation and the environmental implication of its biota at Garyounis, SW Benghazi. The Third International Conference on Basic Sciences & Their Applications. *International Journal of Applied Science*; 2019; Vol 1 (1): 218-235.
15. Muftah AM. and Elebaidi SK. The Geoethical Scenario towards the Archaeological Heritage in Al Jabal Al Akhdar, NE- Libya. *Новые идеи в науках о Земле*; Moscow, Russia; 2019; 112-115.
16. Shaltami, O. R., Fares, F.F., Errishi, H., El Oshaibi, F. M. and Bustany, I. 2017. Rare Earth Element Geochemistry and Isotope Geochronology of Quaternary Calcarene in the Al Jabal Al Akhdar, NE Libya. In: 10th International Conference on Applied Geochemistry (ICG-10), 21-22 Nov. 2017: Proceeding Book: 50-68. Institute of Geosciences Federal University of Rio Grande do Sul Porto Alegre, Brazil.