

Histological Changes Induced by Organophosphate Chlorpyrifos in Liver Tissue of Adult Male Rabbits

Hanan EL Daffri ¹, Najat M.H. Mohammed ^{2*}, Fatma Amraje ¹, Fyrouz Khaled ³, Alnagdy Mohammad Ali ⁴

¹ Histology Department, Faculty of Medicine Omer AL-Mukhtar University, AL-Beyda-Libya.

² Laboratory Medicine Department, Faculty of Medical Technology Omer AL-Mukhtar University, AL-Beyda-Libya.

³ Chemistry Department, Faculty of Science, Omer AL-Mukhtar University, AL-Beyda-Libya.

⁴ General Nursing Department, Faculty of Nursing, Omer AL-Mukhtar University, AL-Beyda-Libya

Received: 22 / 09 / 2022; Accepted: 04 / 12 / 2022

المخلص:

المقدمة: مادة الكلوربيرفوس هي أحد أهم المبيدات الحشرية التي يكثر استخدامها في حماية المزارع والحظائر الحيوانات لما لها من دور في القضاء على الحشرات ولكن تأثيرها قد يكون ضار على حياة الإنسان لذلك هدفت الدراسة لمعرفة تأثير الكلوربيرفوس على الأنسجة الكبدية للأرانب البالغة على المدى الطويل من التعرض وكذلك معرفة الدور الوقائي لفيتامين هاء في حماية خلايا النسيج.

المواد والطرق: عشرون أرنب بالغ تم تقسيمها إلى أربع مجموعات: -المجموعة الأولى ضابطة تم تجريعها فقط ماء المقطر اما المجموعة الثانية فهي المجموعة المعالجة تم تجريعها 33.3م/كم من الكلوربيرفوس، والمجموعة الثالثة تم تجريعها فيتامين هاء فقط 100م/كم والمجموعة الرابعة تم اعطاءها المبيد مع جرعة فيتامين هاء، جرعة واحدة صباحية لمدة 12 اسبوع، وبعدها تم تخدير الحيوانات بالكلوروفورم وإزالة الأرباب ووضعها في الفورمالين ومن ثم معالجتها وصبغها بصبغة الهيماتوكسيلين والأيويسين وصبغة ماسون تراكروم.

النتائج: التجربة بينت أن المبيد له عدة تأثيرات على نسيج الكبد حيث سبب تغيرات في مكونات السيتوبلازم للخلايا الكبدية وظهور علامات التغيرات الدهنية على النسيج، كذلك سبب احتقان الأوعية الدموية وتوسع جدارها بالإضافة إلى ارتشاح بالخلايا الالتهابية وتبين أنها عند تجريع الحيوانات بمضادات الأكسدة فيتامين (هـ) كان هناك تحسن في تقليل ارتشاح الخلايا الالتهابية وكذلك كمية الاحتقان وأصبحت الخلايا مقاربة لشكل الطبيعي فضلا عن بعض الخلايا المتأثرة. يمكن الاستنتاج أن الكلوربيرفوس له تأثير سمي على النسيج الكبدى ولكن يمكن تخفيف هذا التأثير باستخدام مضادات الأكسدة مثل فيتامين هاء.

الكلمات المفتاحية: الكلوربيرفوس، فيتامين هاء، المبيدات الحشرية، الأنسجة الكبدية الأرانب.

Abstract

Background: Chlorpyrifos (CPF) is one of the most common pesticides that is extensively applied in agriculture and animal houses. Several experimental studies on rats demonstrated that exposure to CPF elicits a group of deleterious effects including hepatic dysfunction, immunological defect and neurotoxicity. Moreover, there is concern regarding their deleterious effects on human life. This study investigated the influence of chlorpyrifos on the hepatic tissue of adult male rabbits in long-term exposure and evaluated the protective role of vitamin E in ameliorating toxic impacts.

Methods: Twenty adult male rabbits were used in the study. Their average weight was (27.6 ± 1.891) Kg. They were divided into four groups, five animals in each. The control group received only distilled water. The CPF-treated group was administered 33.3mg/Kg of insecticide orally by gastric gavage. The vitamin E group received only Vit E; 150mg/kg body weight. Animals in the combination group were treated with CPF and given Vit E at the same time. The treatment was applied once daily in the morning after a food supplement for 12 weeks. Animals were then anesthetized with chloroform, their livers were removed, placed in formalin, processed and stained with hematoxylin and eosin, and with Masson's trichrome

Results: Histopathological changes showed that CPF caused degenerative changes in hepatocytes in terms of vacuolization of cytoplasm, congestion, dilatation of the sinusoidal lumen, and leukocytic infiltration of liver parenchyma, as well as the presence of signs of fibrosis manifested by trichrome stain indicating the chronic effect of pesticides. In the combination group, the majority of hepatocytes presented with normal structures, and the amount of fibrosis was diminished around the portal area. Additionally, congestion appeared to a milder degree than in the experimental group, suggesting the antioxidant role of vitamin E in reducing toxic effects. Conclusion: The study concluded that CPF could cause hepatocellular damage to the rabbits' tissue which can be alleviated by applying antioxidants such as vitamin E.

Keywords: Chlorpyrifos, Vit E, pesticides, rabbit's liver.

*Correspondence: Najat Mohammed.

najat.mohammed@omu.edu.ly

1. INTRODUCTION

Chlorpyrifos (CPF) is one of the most popular types of chlorinated organophosphate pesticides (1) that have an insecticidal effect against pests and is vastly applied in a wide field including agriculture, animal husbandry, gardens, and even households (2). CPF is a cholinesterase inhibitor that is commonly used (3). The mechanism of action of cholinesterase inhibitors is the same in both therapeutic and toxic uses. Inhibition of acetylcholinesterase (ACh E) disturbs the dynamic interaction between acetylcholine synthesis, release, and degradation. This causes an accumulation of synaptic ACh levels with prolonged stimulation of cholinergic receptors on postsynaptic cells. This comparative rise in cholinergic signaling leads to practical signs and symptoms of cholinergic toxicity (4). However, the range of cholinergic signs can be diverse with different inhibitors, the same inhibitor in different classes, or the same inhibitor with different ways of exposure (5). The general dysfunction seems due to inhibition, revealing the primary toxic effect on the endocrine system but it can also affect other organs such as the liver and kidneys (1). Chlorpyrifos is categorized as a moderately toxic pesticide in rats and mice with a lethal toxic dose (LD50) between 50-500 mg/kg body weight. Additionally, it shows moderate toxicity in sheep, guinea pigs and pigs (5, 6).

Several experimental studies on rats demonstrated that exposure to CPF elicits a group of deleterious effects including hepatic dysfunction, immunological defect and neurotoxicity (7). Acute and chronic toxicity of CPF is associated with an increased production of reactive oxygen species (ROS), which has been anticipated as a mechanism to create oxidative stress (8, 9). Antioxidants are molecules that can decrease or stop the oxidation of other molecules by ROS (10). Vitamin E is a non-enzymatic antioxidant, which is a main membrane-bound antioxidant employed by the cell (11). It has been reported that vitamin E has an antioxidant role against lipid peroxidation of cell membranes of mammals and has a role in maintaining the activities of mitochondrial enzymes from ROS (12).

The present study aims to investigate the influence of chlorpyrifos on the hepatic tissue of adult male rabbits on long-term exposure. It is also intended to evaluate the protective role of vitamin E in ameliorating toxic impacts.

The present study aims to investigate the influence of chlorpyrifos on the hepatic tissue of adult male rabbits on long-term exposure. It is also intended to evaluate the protective role of vitamin E in ameliorating toxic impacts.

2. METHODS

Chemicals

Commercial formulation of chlorpyrifos (50% EC) of high purity (500g/L) product of Pesticides & Chemicals Company, China. Vitamin E Tablets 100mg were purchased from Sigma Aldrich, India.

Animals

The study was conducted in the Histology Department at the Faculty of Medicine of Omer AL-Mukhtar University. Twenty adult male New Zealander rabbits were used in this study and their average weight was (27.6 ± 1.891) Kg. These animals were divided into four groups with five animals in each. They were housed in cages in the experimental animal house of the

Chemistry Department of the Science Faculty at Benghazi University and they acclimatized for five days before the experiment under providing standard laboratory conditions with free access to food and water. The cleanliness and hygiene of the rabbits were checked regularly and cages were also cleaned of animal waste daily.

Experimental design

Animals were divided into four groups as follows;

- The control group received only distilled water.
- The CPF-treated group were administrated 33.3mg/Kg of insecticide orally by gastric gavage (13).
- The vitamin E group received only Vit E 100mg/kg body weight.
- The combination group those rabbits were given both CPF (33.3mg/Kg) and Vit E (100mg/Kg).

The treatment was applied once daily in the morning after a food supplement for 12 weeks. At the end of the experiment, animals were anesthetized by chloroform and sacrificed. The liver was removed from each animal and fixed by 10% neutral buffered formalin, washed, dehydrated, cleared, and embedded in paraffin. The prepared sections of 5 μ m thickness were stained by Harris Hematoxyline and Eosin (H&E) and Masson's trichrome for histopathological microscopy examination (14).

3. RESULTS

Histological features of control liver sections of rabbits revealed normal hepatic architectures; the central vein (CV) in the center of the lobule and normal sinusoidal lumens (Fig. 1 B). Hexagonal hepatocytes with clear and well-defined nuclei were regularly arranged in the lobule (Fig. 1 B).




Figure 1: Liver H&E sections of the control group showing: A). (x100) Normal hepatic architecture with the CV in the center of the hepatic lobule and portal area in the corners. Section B (x400) presents normal sinusoidal lumens (L), normal arrangement and structure of hepatocytes (red arrow).

The most frequent histopathological changes in the livers of chlorpyrifos-treated rabbits were cytoplasmic vacuolization of the hepatocytes at the three zones of the liver, which are the first signs of fatty changes (Fig. 2, B), often accompanied by sinusoidal dilatation (Fig. 2, D). Additionally, congestion of the CV has been clearly noticed (Fig. 2 C), with infiltration of mononuclear cells also found especially around CV and PA (Fig. 2 A and C).




Figure 2: Light micrographs of the liver sections of CPF-treated rabbits showing:- A and B mononuclear cell infiltration around PA (black arrow) and hepatocytes with cytoplasmic vacuolations (circle). C and D congestion of the central vein (CV) and sinusoidal dilatation (SD). (Mic Mag x100 (A) x400(B) H&E).




Figure 3: Light micrographs of the livers of rabbits of the combination group showing:- A). Marked reduction in inflammatory cells around CV in liver tissue receiving CPF with Vit E (arrowhead). B). Cytoplasmic vacuolation of hepatocytes was milder than of CPF-treated rabbits without Vit E (circle). (Mic Mag x100 (A) x400(B) H&E).

Microscopic examination of liver tissues of rabbits receiving CPF with Vitamin E shows a reasonable reduction in the inflammatory cells around the CV and PA (Fig. 3A). In addition, the cytoplasmic vacuolation of hepatocytes was milder than that noticed in CPF-treated rabbits (Fig. 3B).

Microscopic examination of liver tissues of rabbits treated with CPF with trichome stain showed collagen deposition around PA, which may indicate an early stage of fibrosis (Fig. 4 B) that is virtually invisible on routine hematoxylin and eosin (H&E) stains. However, in samples from rabbits in the combination group, the amount of collagen deposition reduced around PA as shown in figure 4 C.




Figure 4: Light micrographs of liver stained with Masson trichrome showing:- A). Control section of liver tissue with a normal amount of collagen around portal area B). Livers of chlorpyrifos-treated rabbits demonstrated an increased amount of collagen and congestion around the portal triad (arrow). C). The combination group showed improved tissue in terms of reducing collagen fibers in the portal area by giving antioxidants.

4. DISCUSSION

The main site of metabolism and activation of CPF is the liver, through the cytochrome P450 enzyme (15). Various previous studies have shown pathological alterations in the liver tissue of animals after exposure to CPF (16, 17). Thus, this experiment was focused on determining the toxic effect of CPF on the liver tissue of adult male rabbits.

Our results showed congestion of the central vein and dilation of the sinusoidal lumens in all liver specimens of CPF-treated rabbits. These pathological changes are probably due to the toxic effect of CPF. These results are in accordance with some investigators (18, 19). Additionally, vacuolation in the hepatocytes was the main histopathological change in all liver samples that received CPF (20). This could be due to an alteration in lipid metabolism in the liver. This result agreed with an experiment on male rats after CPF exposure, where they found a significant elevation in plasma levels of cholesterol and triglycerides (21). Similar pathological changes in liver tissue were reported in previous studies on CPF (22, 23). Moreover, the presence of monocyte infiltration due to CPF intoxication around the CV and PA is shown in Figure 2 B and was also highlighted after organophosphate s (OP) exposure in a previous study on male rats (1). A similar finding was presented by EL-Demerdash et al who observed that rats treated with CPF had elevated liver enzymes (AST&ALT) indicating intoxication of hepatic tissue, which subsequently leads to enzyme leakage (24). Thus, the release of enzymes provokes inflammatory reactions and liver necrosis as well. Vitamin E (a-tocopherol) is a lipid-soluble antioxidant (25), which prevents chain initiation and propagation of free radical reaction and lipid peroxidation in the cell membrane (26). It also affects the cellular response to oxidative stress through modulation of the signal-transduction pathway and is considered a membrane stabilizer (27-29). The toxic effects of organophosphates (OP) can be restrained by the oral supplementation of antioxidants such as vitamin E (30). Vitamin E has been shown to have an ameliorating effect against the insecticide's toxicity and plays a role in inhibiting the pathological mechanisms of oxidative stress induced by cypermethrin (31, 32).

The present study found that the severity of hepatocellular damage was minimized in CPF-treated rabbits when they were given vitamin E along with insecticides, indicating that vitamin E ameliorates the toxic effect of CPF. This result is in agreement with the previous work investigating the antioxidant effect of vitamin E against the histopathological changes of CPF treating Kunming mice (33). In addition, other researchers have observed that vitamin E is effective in reducing inflammatory reactions and has a role in the protection of hepatocytes from oxidative stress injury induced by OP in rat tissue (12, 34). Moreover, vitamin E has had an effective role in the prevention of oxidative damage to the lung tissue of CPF-treated animals (35).

In our study, Masson's trichrome stain revealed moderate liver fibrosis between hepatic lobules and around the portal area in the CPF-treated group as shown in Figure 4C; a similar observation has been reported by previous findings (36). This fibrosis is considered the end-stage hepatic injury that usually appeared upon chronic exposure and is consistent with our experiment period. The mechanism of developing liver fibrosis is sophisticated but it is mainly participated by increased reactive oxygen species (ROS) and inhibiting antioxidants of the

biological system (35, 37). Interestingly, in this experiment, the amount of fibrosis was reduced in rabbit tissue by giving vitamin E with insecticides as compared with the CPF-treated group.

5. CONCLUSION:

It can be concluded that CPF could have a destructive effect on liver tissue and has the risk of being harmful to human life. However, this hepatic damage can be modulated and diminished by using vitamin E as dietary supplementation.

Study limitations

A possible limitation of the current research study is the absence of serum liver enzymes that could have contributed to the valuation of functional liver involvement in CPF intoxication.

6. REFERENCES:

1. Cobilinschi C, Țincu R, Neagu T, Becheanu G, Sinescu R, Checheriță I, et al. Histopathological features of low-dose organophosphate exposure. *Rom J Morphol Embryol*. 2020;61(2):423-32.
2. Ma P, Wu Y, Zeng Q, Gan Y, Chen J, Ye X, et al. Oxidative damage induced by chlorpyrifos in the hepatic and renal tissue of Kunming mice and the antioxidant role of vitamin E. *Food and chemical toxicology*. 2013(58):177-83.
3. Bartles E, Nachmansohn D. Organophosphate inhibitors of acetylcholine-receptor and -esterase tested on the electroplax. *Arch Biochem Biophys*. 1969;133(1):1-10.
4. Gultekin F, Delibas N, Yasar S, Kilinc I. In vivo changes in antioxidant systems and protective role of melatonin and a combination of vitamin C and vitamin E on oxidative damage in erythrocytes induced by chlorpyrifos-ethyl in rats. *Arch Toxicol*. 2001; 75(2):88-96.
5. Pope C, Karanth S, Liu J. Pharmacology and toxicology of cholinesterase inhibitors: uses and misuses of a common mechanism of action. *Environmental Toxicology and Pharmacology* 2005;19 (3):433-46.
6. Ishii S, Bell J N, Marshall F M. Phytotoxic risk assessment of ambient air pollution on agricultural crops in Selangor State, Malaysia. *Environmental Pollution*. 2007;150(2):267-79.
7. Babazadeh M, Najafi G. Effect of chlorpyrifos on sperm characteristics and testicular tissue changes in adult male rats. *Veterinary Research Forum* 2017;8(4):319 - 26.
8. Bagchi D, Bagchi M, Hassoun E A, Stohs SJ. In vitro and in vivo generation of reactive oxygen species, DNA damage and lactate dehydrogenase leakage by selected pesticides. *Toxicology and Applied Pharmacology*. 1965;104(1-3):129-40.
9. Chargui I, Grissa I, Bensassi F, Hrira MY, Haouem S, Haouas Z, et al. Oxidative Stress, Biochemical and Histopathological Alterations in the Liver and Kidney of Female Rats Exposed to Low Doses of Deltamethrin (DM): AMolecular Assessment. *Biomed Environ Sci*. 2012; 25(6):672-83.
10. Owuor ED, Kong ANT. Antioxidants and oxidants regulated signal transduction pathways. *Biochem Pharmacol*. 2002;64(5-6):765-70.
11. Burton GW, Ingold KU. Vitamin E as an in vitro and in vivo antioxidant. *Ann NY Acad Sci*. 1989;570:7-22.
12. Amin KA, Hashem KS. Deltamethrin-induced oxidative stress and biochemical changes in tissues and blood of catfish (*Clarias gariepinus*): antioxidant defense and role of alpha-tocopherol. *BMC Veterinary Research* 2012;8(1):1-8.
13. Yassin MM, Adas TO, Yasin MM. Serum Glucose, Bilirubin, Liver Enzymes, Renal Parameters, Protein Profile and Some Electrolytes in Adult Male Domestic Rabbits Intoxicated with Chlorpyrifos. *J Toxicol Risk Asses*. 2021;7(1):037.
14. Bancroft JD, Gamble M. Theory and practice of histological techniques. 6th ed. Philadelphia: Churchill Livingstone Elsevier; 2008. 126-7 p.
15. Mutch E, Williams FM. chlorpyrifos and parathion are metabolised by multiple cytochromes P450 in human liver. *Toxicology* 2006;224(1-2):22-32.
16. Tanvir E, Afroz R, Chowdhury M A, Gan S H, Karim N, et al. A model of chlorpyrifos distribution and its biochemical effects on the liver and kidneys of rats. *Human and Experimental Toxicology*. 2016;35(9):991-1004.
17. El durssi I, Elawami I, Mahmoud G, Benkhaia F, Mohamed N. Experimental Studies on the effects of Chlorpyrifos on Rats, Histopathological changes. *Mukhtar Journal*. 2006;(13):67-79.
18. Ezzi L, Belhadj S I, Haouas Z, Sakly A, Grissa I, Chakroun S. Histopathological and genotoxic effects of chlorpyrifos in rats. *Environmental Science and Pollution Research*. 2016;23(5):4859-67.
19. Tripathi S, Srivastav A K. Liver profile of rats after long-term ingestion of different doses of chlorpyrifos. *Pesticide biochemistry and physiology*. 2010; 97(1):60-5.
20. Devi Y, Mishrab A. Histopathological Alterations in Gill and Liver Anatomy of Fresh Water, Air Breathing Fish *Channa Punctatus* after Pesticide Hilban (Chlorpyrifos) Treatment. *Advances in Bioresearch*. 2013;4(2):57- 62.
21. Slotkin TA, Brown K K, Seidler F J. Developmental exposure of rats to chlorpyrifos elicits sex-selective hyperlipidemia and hyperinsulinemia in adulthood. *Environmental health perspectives*. 2005;113(10):1291-4.
22. Heikal T, Mossa A, Marei G, Rasoul M. A. Cyromazine and chlorpyrifos induced renal toxicity in rats: the ameliorating effects of green tea extract. *J Environ Anal Toxicol*. 2012; 2(5).
23. Tanvir EM, Afroz R, Chowdhury M, Khalil M, Hossain M, Rahman MA. Honey has a protective effect against chlorpyrifos-induced toxicity on lipid peroxidation, diagnostic markers and hepatic histoarchitecture *European Journal of Integrative Medicine*. 2015;7(5):525-33.

24. El-Demerdash FM. Antioxidant effect of vitamin E and selenium on lipid peroxidation, enzyme activities and biochemical parameters in rats exposed to aluminium. *Journal of Trace Elements in Medicine and Biology*. 2004;18(1):113-21.
25. Machlin LJ. Vitamin E, a comprehensive treatise. . Marcel Dekker.1980.
26. Kamal- Eldin A, Appelqvist L. The chemistry and antioxidant properties of tocopherols and tocotrienols. *J Lipids*. 1996; 31(7):671-701.
27. Azzi A, Boscoboinik D, Hensey C. The protein kinase C family. *European Journal of Biochemistry*. 1992;208(3):547-57.
28. Traber MG, Packer L. Vitamin E: beyond antioxidant function. *The American journal of clinical nutrition* 1995;62(6):1501s-9s.
29. Clarke MW, Burnett JR, Croft K. Vitamin E in human health and disease. *Critical reviews in clinical laboratory sciences*. 2008;45(5):417-50.
30. Ahmad LA, Khan A, Khan MZ. Cypermethrin induced biochemical and hepato-renal pathological changes in rabbits. *Int J Agr Biol* 2012;13(6):865-72.
31. Raina R, Verma PK, Pankaj NK, Kant V. Ameliorative effects of alpha-tocopherol on cypermethrin induced oxidative stress and lipid peroxidation in Wistar rats. *Int J Med Sci*. 2009;1:396-9.
32. Sallam M A, Ahmad M, Ahmad I, Gul S T, Idrees M, Bashir M I, et al. Toxic Effects of Cypermethrin on the Reproductive Functions of Female Rabbits and Their Amelioration with Vitamin E and Selenium. *Pak Vet J*. 2015;35(2):193-6.
33. Ma P, Wu Y, Zeng Q, Gan Y, Chen J, Ye X, & Yang X. Oxidative damage induced by chlorpyrifos in the hepatic and renal tissue of Kunming mice and the antioxidant role of vitamin E. *Food and chemical toxicology*. 2013;58,:177-83.
34. Al-Omar M S, Naz M, Mohammed S, Mansha M, Ansari M, Rehman N, et al. Pyrethroid-Induced Organ Toxicity and Anti-Oxidant-Supplemented Amelioration of Toxicity and Organ Damage: The Protective Roles of Ascorbic Acid and α -Tocopherol. *Int J Environ Res Public Health*. 2020;17:1-27.
35. Hassani S, Sepand MR, Jafari A, Jaafari J, Rezaee R, Zeinali M, et al. Protective effects of curcumin and vitamin E against chlorpyrifos-induced lung oxidative damage *Hum Exp Toxicol*. 2015; 34(6): 668-76.
36. Laothong U. Pesticides and Non-alcoholic Fatty Liver Disease. *Thia J Toxicol*. 2017; 31 (2):97-107.
37. Li S, Zheng X, Zhang X, Yu H, Hana B, Lv Y, et al. Exploring the liver fibrosis induced by deltamethrin exposure in quails and elucidating the protective mechanism of resveratrol. *Ecotoxicol Enviro Saf*. 2021;207:111501.