

Effects of Acetaminophen on the Reproductive Parameters and the Ameliorative Effects of Rutin and Mesenchymal Stem Cells in Male Rats

Rabihah E. A. Elduob¹, Eda M. A. Alshailabi^{1*}, Samia M. Efkeren¹

¹ Zoology Department- Science Faculty- Omar Al-Mukhtar University.

Received 16/ 10 / 2023; Accepted 15 / 10 / 2023

المخلص

صممت الدراسة الحالية لتقييم آثار الأسيتامينوفين على العوامل الإنجابية والتأثيرات التحسينية للروتين والخلايا الجذعية الوسيطة في ذكور الجرذان. وقد تم تقسيمهم إلى دراستين، أولاً: تم استخدام عشرين جرذاً صغيراً كمصدر للخلايا الجذعية المشتقة من نخاع العظم، ثانياً: تم تقسيم خمسين من ذكور الجرذان البالغة إلى 5 مجاميع: المجموعة الضابطة (G1)، وأعطيت الجرذان الأسيتامينوفين (750 ملجم/كجم من وزن الجسم) كل 72 ساعة لمدة 21 يوماً، ثم تركت لمدة 30 و60 يوماً دون معاملة (G2)، أعطيت الجرذان الأسيتامينوفين لمدة 21 يوماً ثم عولجت بروتين (25 ملجم/كجم من وزن الجسم) لمدة 30 و60 يوماً (G3)، تم إعطاء الجرذان الأسيتامينوفين لمدة 21 يوماً، وتم حقنها بواسطة خلايا جذعية (1.5×10^6 خلية في 0.5 من محلول فوسفات البوتاسيوم) في الوريد الذيلي لمدة 30 و60 يوماً (G4)، وتم إعطاء الجرذان الأسيتامينوفين لمدة 21 يوماً، وحقنها بواسطة الخلايا الجذعية في الوريد الذيلي ثم عولجت بالروتين لمدة 30 و60 يوماً (G5). أنتج إعطاء الأسيتامينوفين انخفاضاً ملحوظاً في القيمة المتوسطة لمستويات هرمونات التستوستيرون، والملوتن، والهرمون المنبه للجريب بعد 30 و60 يوماً مقارنةً بالمجموعة الضابطة. حيث أن الجرذان المعالجة بالروتين والخلايا الجذعية معاً أظهرت زيادة كبيرة في متوسط قيم مستوى هذه الهرمونات بعد 30 و60 يوماً بالمقارنة مع مجموعة الأسيتامينوفين. وفي الختام، أظهرت هذه الدراسة أن الروتين والخلايا الجذعية عند إعطائهما معاً يحميان من الانخفاض الناجم عن الأسيتامينوفين في مستوى هرمونات الخصية.

الكلمات المفتاحية: وظائف الخصية، روتين، خلايا اللحمية المتوسطة، أسيتامينوفين، الجرذان.

Abstract

The present study has been designed to evaluate the effects of acetaminophen on the reproductive parameters and the ameliorative effects of rutin and Mesenchymal stem cells in male rats. The rats were divided into two studies: firstly, twenty young male rats were used as a source of bone marrow-derived MSCs. Secondly, fifty adult male rats were divided into 5 groups: Control group (G1); rats were administrated of AAP (750 mg/kg b.w.) every 72h for 21 days, then left for 30 and 60 days without treatment (G2), rats were administrated of AAP for 21 days then treated with RT (25mg/kg b.w/d) for 30 and 60 days (G3), rats were administrated of AAP for 21 days then the rats were injected by BM-MSCs (1.5×10^6 cells in 0.5 PBS) in the tail vein for 30 and 60 days (G4), and rats were administrated of AAP for 21 days then the rats were injected by BM-MSCs in the tail vein, then treated with RT for 30 and 60 days (G5). Administration of AAP produced a significant decrease in the mean value of the Te, LH, and FSH levels after 30 and 60 days as compared to the C groups. Whereas rats treated with RT and MSCs showed a significantly high increase in the mean values of these hormone levels after 30 and 60 days compared with the AAP group. In conclusion, this study demonstrated that RT and MSCs when treated in combination, are protected against the AAP-induced decrease in the hormone level of testicular.

Keywords: testes functions; rutin; mesenchymal cells; acetaminophen; rats.

1. INTRODUCTION

The properties of medicinal plants are mainly due to the presence of various complex chemical substances of different composition which occur as secondary metabolites [1]. Rutin (RT) has several pharmacological properties, including antioxidant, anticarcinogenic, cytoprotective, vasoprotective, cardioprotective and neuroprotective activities [2, 3]. Moreover, RT is a flavonoid of the flavonol type that is found in many typical plants, such as buckwheat, passion flower, apple and tea. It is also an important dietary constituent of foods and plant-based beverages [4, 5].

On the other hand, RT is known as Vitamin P, and it has been extensively studied and known to exhibit multiple pharmacological activities including antibacterial and antiviral, antiprotozoal, antitumor, antiallergic, anti-inflammatory, and antiplatelets [6, 7].

Acetaminophen (AAP) is the drug of choice for patients who cannot use nonsteroidal anti-inflammatory drugs (NSAIDs) because of hypersensitivity to aspirin, gastric ulcers, impaired blood coagulation, pregnancy, breastfeeding, or fever accompanying disease in children [8]. AAP, also known as (paracetamol or N-acetyl-p-amino-phenol) is the most commonly used antipyretic and pain reliever, and since 1955 it has been a widely analgesic medication in many countries [9,10]. Moreover, AAP is metabolized in the liver by cytochrome P450 (CYP450) enzymes, to N-acetyl-p-benzoquinone imine

*Correspondence: Eda M. A. Alshailabi

eda.muftah@omu.edu.ly

(NAPQI). NAPQI reacts with glutathione (GSH), therefore, overdoses of paracetamol may result in a depletion of hepatocellular GSH [11, 12]. Acute overdose of AAP might cause toxicity of testis in humans and experimental animals [13]. Furthermore, the long-term use of AAP causes toxicity effects in the organs, including hepatotoxicity, renal toxicity, and testicular toxicity. In addition, it affects the blood chemistry and reproductive parameters, [14, 15] such as semen quality, particularly sperm morphology and its fertilizing ability [16]. Also, [14] it was found that the long-term AAP caused an increased risk of oxidative stress, testicular tubules, blood cells, and causing hypertension and heart infarction. However, the administration of AAP is one of the most common causes of toxicity worldwide [17]. Moreover, the AAP-toxicity is the production of the reactive intermediate NAPQI by CYP450 which is removed by conjugation to glutathione in the therapeutic doses, so the higher doses of AAP lead to the drop of cellular content GSH which allows NAPQI to combine with the cellular proteins and induction of lipid peroxidation, leading to toxicity [18, 13, 19].

Bone marrow mesenchymal stem cells (MSCs) are a type of mesoderm-derived stem cells that can differentiate into a variety of cell types, including osteoblasts, chondrocytes, adipocytes, and muscle cells [20], where MSCs have the ability to differentiate into germ cells [21]. Moreover, MSC-secreted factors may protect the spermatogenic dysfunction after busulfan treatment by reducing the apoptosis of spermatogenic cells and enhancing the expression of intercellular adhesion molecules [22]. MSCs are used in the repair and reconstruction of tissues and organs, treatment of degenerative diseases, and anti-aging treatment due to their self-renewal and multidirectional differentiation potential [23, 24].

So, this study aimed to evaluate the effects of acetaminophen on the reproductive parameters and the ameliorative effects of rutin and Mesenchymal stem cells in male rats.

2. MATERIALS AND METHODS:

Chemicals:

- Acetaminophen ($C_8H_9NO_2$) was used. It was purchased from Sigma Chemical Company (USA) [25].
- Rutin ($C_{27}H_{30}O_{16}$) the natural antioxidant was purchased from Sigma Chemical Company (USA) [26].
- Bone marrow-derived stem cells, one important source of mesenchymal stem cells (MSCs), have been isolated and cultured at the Medical Research Center, Aleibbasiuh, Ain Shams University.

Experimental Animals:

A total number of 70 male albino rats (*Rattus norvegicus*) were used. They were divided into two main studies: (S1) consisted of twenty young male albino rats of average weight 100 g that were used as a source of bone marrow-derived MSCs, and (S2) consisted of fifty adult male albino rats that were divided into 5 groups with an average weight of (150-160g) for the experimental study. The rats were obtained from Animal House of El-Salam Farm, Giza-Cairo, Egypt, and were acclimatized to the laboratory conditions for two weeks prior to the start of the experiment. They were housed in metabolic

cages at a temperature of 24-27 °C, 12 hours dark/ light cycle, and received standard food and water ad-libitum with fresh daily supplies. The experimental procedures complied with the guidelines of the Committee on Care and Use of Experimental Animal Resources, Ain Shams University, Cairo, Egypt.

Experimental Design:

The first study (S1): Preparation of Bone Marrow-derived Mesenchymal Stem Cells (MSCs):

Twenty young male albino rats of average weight 100 g (6 weeks old) were used as a source of bone marrow-derived MSCs [27]. The rats were injected with BM-MSCs (1.5×10^6 cells in 0.5 PBS) [28] in the tail vein [29]. The cultured BM-MSCs were characterized by using NAVIOS flow cytometer by BECKMAN COULTER in the Medical Research Center of Ain Shams University [30].

The second study (S2):

Fifty adult male rats were randomized into five groups 10 rats in each:

Group (1): In the control group (C), the rats were left as the normal control rats with no treatment.

Group (2): Acetaminophen treated group (AAP); the rats were orally administrated with a dose of AAP (750 mg/kg b.w./ every 72h) for 21 days. Then, they were left for 30 and 60 days without any treatment.

Group (3): Acetaminophen with rutin group (AAP+RT), the rats received oral doses of AAP (750 mg/kg b.w./ every 72h) for 21 days. Then, they were treated orally with RT at a dose of (25mg/kg b.w./d) for 30 and 60 days.

Group (4): Acetaminophen with stem cells group (AAP+MSCs), the rats received oral doses of AAP (750 mg/kg b.w./ every 72h) for 21 days. Then, the rats were injected with BM-MSCs (1.5×10^6 cells in 0.5 PBS) in the tail vein for 30 and 60 days.

Group (5): Acetaminophen with stem cells and rutin group (AAP+MSCs+RT), the rats received oral doses of AAP (750 mg/kg b.w./ every 72h) for 21 days. Then, the rats were injected with BM-MSCs (1.5×10^6 cells in 0.5 PBS) in the tail vein with treated orally by rutin at a dose of (25mg/kg b.w./d) for 30 and 60 days.

At the end of the experimental period, the rats were overnight fasted after the last dose and blood samples were collected.

Determination of Serum Hormone Levels:

Blood samples were taken into clean and dry screw-capped centrifuge tubes and then centrifuged at 3000rpm for 15 minutes in order to separate clear serum samples. They were then stored at -20°C until used for the determination of different biochemical parameters. Sera were used for the determination of male sex hormones analysis such as the testosterone hormone (Te), luteinizing hormone (LH), and follicle-stimulating hormone (FSH), were evaluated to determine the sex hormones activities of the control group, and the experimental groups were estimated according to the method of [31] for Te, and [32] for FSH [33] for LH.

Statistical Analysis:

Results were expressed as mean standard error (SE). The parameters were analyzed using significance by one-way ANOVA. Means were detached using Tukey's test at $P < 0.05$. Also, using the T-test to compare between two means. All statistical procedures were performed with (Minitab version 17).

3. RESULTS: -

Determination of the Testosterone Hormone Level (Te): -

Averages of the Te level were given in Table (1), it has shown a significant decrease ($P < 0.05$) in the mean value of the Te level after 30 and 60 days in the AAP groups (0.8200 ± 0.37 & 1.2000 ± 0.07) as compared to the C group (4.340 ± 0.246 & 4.340 ± 0.25). Whereas there was no significant ($P < 0.05$) between the AAP and AAP+RT groups (1.2000 ± 0.031 & 1.6600 ± 0.051) in the mean value of the Te level after 30 and 60 days. Also, the mean values of the Te level after 30 and 60 days showed a significant increase in the treated group with the AAP+MSCs groups (1.8200 ± 0.058 & 2.600 ± 0.0511) when compared with the AAP groups. Moreover, the treated groups by AAP+MSCs+RT in the mean value of the Te level after 30 and 60 days showed a significantly high increase (2.5000 ± 0.044 & 2.820 ± 0.153) when compared with AAP groups.

Determination of the Luteinizing Hormone (LH) Level: -

Data recorded for the mean values of the LH level of control and experimental rats were obtainable in Table (1), a significant decline ($P < 0.05$) was found in the mean value of LH level after 30 and 60 days in the AAP rats (0.5200 ± 0.037 & 1.2600 ± 0.075) as compared to the C rats (5.10 ± 1.358 &

5.10 ± 0.23). Although, there was a significant increase ($P < 0.05$) in the AAP+RT group (1.3000 ± 0.031) in the mean value of LH level after 30 days when compared to the AAP rats (0.5200 ± 0.037). No significant effects were observed on the AAP+RT rats (2.1400 ± 0.051) in the mean value of LH level after 60 days as compared to the AAP rats (1.2600 ± 0.075). In addition, the mean values of the LH level after 30 and 60 days revealed a significant increase in the AAP+MSCs groups (1.6800 ± 0.037 & 2.680 ± 0.455) compared to the AAP rats in the mean value of LH level after 30 and 60 days. Furthermore, the mean value of LH level after 30 and 60 days demonstrated a significantly high increase in the AAP+MSCs+RT rats (2.9200 ± 0.066 & 4.180 ± 0.199) when compared with AAP groups.

Determination of the Follicle-stimulating Hormone (FSH) Level: -

On measuring the FSH level, the data presented in Table (1) showed a significant decrease ($P < 0.05$) in the mean value of the FSH level in the AAP groups after 30 and 60 days (0.6400 ± 0.040 & 1.4000 ± 0.070) compared to the C group (4.560 ± 0.213 & 4.60 ± 0.21). Additionally, the mean value of FSH level showed a significant increase in the AAP+RT group (1.5600 ± 0.087) after 30 days when compared with the AAP rats (0.6400 ± 0.040). Moreover, there were no remarkable changes between the mean value of the FSH level after 60 days in the AAP group (1.4000 ± 0.070) and the AAP+RT group (3.2200 ± 0.20). While the AAP+MSCs rats (1.8400 ± 0.024 & 3.200 ± 0.170) presented a significant increase ($P < 0.05$) in the mean value of the FSH level after 30 and 60 days when compared with the AAP groups. However, the mean values of the FSH level after 30 and 60 days revealed a notably high increase in the AAP+MSCs+RT groups (2.680 ± 0.120 & 4.640 ± 0.301) when compared with AAP groups.

Table 1: Average mean values of creatinine, urea and albumin levels in control and experimental groups.

Duration	Parameter	C	AAP	AAP + RT	AAP + MSCs	AAP+ MSCs+ RT
30 days	Te (ng/ml)	4.340 ± 0.246^A	0.8200 ± 0.37^D	1.2000 ± 0.031^D	1.8200 ± 0.058^B	2.5000 ± 0.044^C
	LH (mIU/ml)	5.100 ± 0.219^A	0.5200 ± 0.037^D	1.3000 ± 0.031^C	1.6800 ± 0.037^C	2.9200 ± 0.066^B
	FSH (mIU/ml)	4.560 ± 0.213^A	0.6400 ± 0.040^D	1.5600 ± 0.087^C	1.8400 ± 0.024^C	2.680 ± 0.120^B
60 days	Te (ng/ml)	4.340 ± 0.25^A	1.2000 ± 0.07^D	1.6600 ± 0.051^{CD}	2.600 ± 0.0511^C	2.820 ± 0.153^B
	LH (mIU/ml)	5.10 ± 0.23^A	1.2600 ± 0.075^C	2.1400 ± 0.051^{BC}	2.680 ± 0.455^B	4.180 ± 0.199^A
	FSH (mIU/ml)	4.60 ± 0.21^A	1.4000 ± 0.070^C	2.1000 ± 0.083^C	3.200 ± 0.170^B	4.640 ± 0.301^A

Data are expressed as mean $\pm$ SE of rat within each row, means with different superscript (A, B, C & D) were significantly different at $P < 0.05$, were means superscripts with the same letters mean that there is no significant difference ($P < 0.05$).

*C= Control group, AAP= Acetaminophen treated group, AAP+RT= Acetaminophen with rutin group, AAP + MSCs = Acetaminophen with stem cells group, and AAP+MSCs +RT = Acetaminophen with stem cells and rutin group.

4. DISCUSSION:

The results showed that AAP produced a significant decrease ($P < 0.05$) in the mean value of the Te, LH, and FSH levels after 30 and 60 days as compared to the C group. Similar results were obtained by [13, 17] who found that the administration of AAP revealed a significant decrease in reproductive parameters. A significant reduction in the Te, LH, and FSH levels in treated groups by AAP indicates that the AAP may cause testicular toxicity and impaired fertility. Another study investigated the toxic effects of a high dose of AAP on the reproductive system of male rabbits and mice, which induced several changes and harmfully affected the histological structure of the seminiferous tubules [34, 35, 13]. So, the results suggested that AAP can potentially cause reproductive toxicity when high and long-term treatment.

The AAP+MSCs rats obtained a significant increase ($P < 0.05$) in the mean value of the Te, LH, and FSH levels after 30 and 60 days when compared with the AAP rats. These are in agreement with the results of [36, 37]. These effects may be due to MSCs differentiation into germ cells, where the mechanisms involved in the restorative effects of MSCs on testicular functions are due to the ability of transplanted MSCs to stimulate the production of substances capable of inhibiting ROS, cell death, inflammation, and mutagenic activities [36]. On the other hand, MSCs play a role in restoring spermatogenesis by differentiating into sperm cells or maintaining spermatogonial stromal cells [38]. Hence, MSCs could be an essential and powerful approach to treating infertility [36].

In the present study, animals treated with both MSCs and RT after being treated by AAP showed a significantly high increase in the mean values of the Te, LH, and FSH levels after 30 and 60 days were markedly improved when compared with the AAP alone. Combined administration of MSCs and RT reversed the levels of these hormones almost back to nearly normal. Many studies recorded that MSCs or RT treatment resulted in a rapid reversal of testicular injury and marked restoration of normal histological structure of testes [38, 39, 36, 40]. RT administration is an important defensive antioxidant enzyme in tissues, and it preserves the morphology of the testis and spermatogenesis. Moreover, RT is a strong antioxidant and has potent pharmacological capabilities including, anti-inflammatory, immunomodulatory antiviral, anti-angiogenic, anti-mutagenic, and antidiarrheal [40]. Also, [41] reported that the RT detoxifies the oxidative stress produced in the body by various drugs and chemicals, and showed RT has inhibitory effects against the generation of ROS and membrane lipid peroxidation. On the other hand, [42] it demonstrated that the specific responses of MSCs to oxidative stress may play a crucial role in the regulation of tissue homeostasis as well as

regeneration of organs after oxidative damage. Also, they said, that the MSCs might have a potential role in treating male infertility and testosterone deficiency. Furthermore, Wang et al. [21] found that MSCs can regulate testicular and ovarian function.

5. CONCLUSION:

In conclusion, this study demonstrated that RT and MSCs when administered in combination, protected against the AAP-induced decrease in the hormone level of testicular, suggesting that RT and MSCs exhibited anti-inflammatory potencies against inflammatory toxicities and testicular function changes induced by AAP.

6. REFERENCES:

1. Dar AI, Saxena RC, Bnsal SK. Assessment of hepatoprotective activity of fruit pulp of *Feronia limonia* (Linn.) against paracetamol induced hepatotoxicity in albino rats. Nat. Prod. Plant Resour. 2012, 2 (2):226-233.
2. Javed H, Khan MM, Ahmad A, Vaibhav K, Ahmad ME, Khan A, Ashafaq M, Islam F, Siddiqui MS, Safhi MM, Islam F. Rutin prevents cognitive impairments by ameliorating oxidative stress and neuroinflammation in rat model of sporadic dementia of Alzheimer type. Neuroscience. 2012, 17:340–352.
3. Prasad R, Prasad SB. Antitumor activity of rutin-cisplatin in combination and its protective effect against hematotoxicity. RJLBPCS. 2018, 4(6): 42-47. DOI: 10.26479/2018.0406.04.
4. Kuntić V, Pejić N, Ivković B, Vujić Z, Ilić K, Mičić S, Vukojević V. Isocratic RP-HPLC method for rutin determination in solid oral dosage forms. J Pharm Biomed Anal. 2007, 43:718–721. DOI: 10.1016/j.jpba.2006.07.019.
5. Motamedshariaty VS, Farzad SA, Nassiri-Asl M, Hosseinzadeh H, Effects of rutin on acrylamide-induced neurotoxicity. DARU J Pharmaceutical Sci. 2014, 22 (27): 1-9. Doi: 10.1186/2008-2231-22-27.
6. Ganeshpurkara A, Salujaa AK. The Pharmacological Potential of Rutin. Saudi Pharm J. 2017, 25(2): 149–164. Doi: 10.1016/j.jsps.2016.04.025.
7. Toro RM, Aragóna DM, Ospinaa LF, Ramosb FA, Castellanosb L. Phytochemical Analysis, Antioxidant and Anti-Inflammatory Activity of Calyces from *Physalis peruviana*. NPC. 2014, 9 (11): 1573 –1575.
8. Benista MJ, Nowak JZ. Paracetamol: Mechanism of action, applications and safety concern. Acta Pol. Pharm. Drug. Res. 2014, 71: 11–23.
9. Hasanein P, Sharifi M. Effects of rosmarinic acid on acetaminophen-induced hepatotoxicity in male Wistar rats. Pharm. Biol. 2017, 55(1):1809- 1816. DOI: 10.1080/13880209.2017.1331248.
10. Tittarelli R, Pellegrini M, Scarpellini MG, Marinelli E, Bruti V, Di luca NM, Busardò FP, Zaami S.

- Hepatotoxicity of paracetamol and related fatalities. Eur. Rev. Med. Pharmacol. Sci. 2017, 21 (1): 95-101.
11. Sabiu S, Sunmonu TO, Ajania EO, Ajiboye TO. Combined administration of silymarin and vitamin C stalls acetaminophen-mediated hepatic oxidative insults in Wistar rats. Rev. Bras. Farmacogn. 2015, 25: 29-34. DOI: 10.1016/j.bjp.2014.11.012.
 12. Elkomy A, Aboubakr M, Ashraf L. Ameliorative effect of thymus oil on paracetamol induced hepato-renal toxicity: A biochemical, antioxidant and histopathological studies. J. Pharmacol. Clin. Res. 2017, 4(3): 1-8. DOI:10.19080/JPCR.2017.04.555637.
 13. Mohammed HU, Sabry RM. The possible role of curcumin against changes caused by paracetamol in testis of adult albino rat (histological, immunohistochemical and biochemical study). Personal non-commercial use only. EJH copyright. 2020, 43 (3): 819-834. DOI: 10.21608/ejh.2019.18599.1189.
 14. Olaleye MT, Rocha BTJ. Acetaminophen induced liver damage in mice: Effect of some medicinal plants on the oxidative defense system. Exp. Toxicol. Pathol. 2008, 59: 319–327. DOI: 10.1016/j.etp.2007.10.003.
 15. Khayyat LI. Evaluate of the efficacy of Extra *Virgin olive* oil against paracetamol overdose induced renal toxicity in albino rats. Umm. Al-Qura Univ. J. App. Sci. 2018, 5(1): 1-13.
 16. Khayyat LI. Extra *Virgin olive* oil protects the testis and blood from the toxicity of paracetamol (overdose) in adult male rats. Biology 2021, 10: 1042. DOI:10.3390/biology10101042.
 17. Yoon E, Babar A, Choudhary M, Kutner M, Pyrsopoulos N. Acetaminophen-induced hepatotoxicity: A comprehensive update. JCTH. 2016, 4:131–142 DOI: [10.14218/JCTH.2015.00052](https://doi.org/10.14218/JCTH.2015.00052).
 18. Matic MM, Milošević MD, Milica G. Paunović MD, Ognjanović BL, Štajn AS, Saičić ZS. Paracetamol-induced changes of haematobiochemical and oxidative stress parameters in rat blood: Protective role of Vitamin C and β -glucan. Kragujevac J. Sci. 2016, 38: 135-146. DOI: 591.111.1:615.212.3:598.323.4.
 19. Alshailabi EMA, Abdalally, OA, Majeed SF. Histopathological study on the protective effect of vitamin C against paracetamol-induced acute hepatic damage in rat. Global Libyan J. 2021, 53: 1-15.
 20. Sheng G. The developmental basis of mesenchymal stem/stromal cells (MSCs). BMC Developmental Bio. 2015, 15(44): 1-8. DOI 10.1186/s12861-015-0094-5.
 21. Wang Z, Yanga T, Liua S, Chen Y. Effects of bone marrow mesenchymal stem cells on ovarian and testicular function in aging Sprague-Dawley rats induced by D-galactose CELL CYCLE. 2020, 19 (18): 2340–2350. DOI:10.1080/15384101.2020.1806434.
 22. Ghasemzadeh-Hasankolaie M, Eslaminejad MB, Sedighi-Gilani M. Derivation of male germ cells from ram bone marrow mesenchymal stem cells by three different methods and evaluation of their fate after transplantation into the testis. In Vitro Cell Dev Biol Anim. 2016, 52: 49–61. DOI:10.1007/s11626-015-9945-4.
 23. Broekman W, Amatngalim GD, de Mooij-eijk Y, Oostendorp J, Roelofs H, Taube C, Stolk J, Hiemstra PS. TNF- α and IL-1 β -activated human mesenchymal stromal cells increase airway epithelial wound healing in vitro via activation of the epidermal growth factor receptor. Respir Res. 2016;17(3): 1-12. DOI:10.1186/s12931-015-0316-1.
 24. Spitzhorn LS, Megges M, Wruck W, Rahman MS, Otte J, Degistirici O, Meisel R, Sorg RV, Oreffo ROC, Adjaye J. Human iPSC-derived MSCs (iMSCs) from aged individuals acquire a rejuvenation signature. Stem Cell Res Ther. 2019, 10 (100). DOI:10.1186/s13287-019-1209-x.
 25. Anbarasu C, Rajkapoor B, Kalpana J. Protective effect of *Pisonia aculeata* on paracetamol induced hepatotoxicity in rats. J. Exp. Int. Med. 2011, 1(3):167-172.
 26. Shenbagam M, Nalini N. Dose response effect of rutin a dietary antioxidant on alcohol-induced prooxidant and antioxidant imbalance-a histopathologic study. Fund. Clin. Pharm. 2011, 25: 493–502. DOI: [10.1111/j.1472-8206.2010.00861.x](https://doi.org/10.1111/j.1472-8206.2010.00861.x).
 27. Nagaya N, Fujii T, Iwase T, Ohgushi H, Itoh T, Uematsu M, Yamagishi M, Mori H, Kangawa K, Kitamura S. Intravenous administration of mesenchymal stem cells improves cardiac function in rats with acute myocardial infarction through angiogenesis and myogenesis. Am J Physiol Heart Circ Physiol. 2004, 287: 2670 –2676. DOI:10.1152/ajpheart.01071.2003.
 28. Oskoue BN, Lamirault AG, Joseph BC, Treuer AAV, Landa AS, Da Silva AJ, Hatzistergos AK, Dauer M, Balkan AW, Mcniece AI, Harea AJM. Increased potency of cardiac stem cells compared with bone marrow mesenchymal stem cells in cardiac repair. Stem Cells Transl Med. 2012, 1(2): 116–124, DOI: 10.5966/sctm.2011-0015.
 29. Jiang W, Ma A, Wang T, Han K, Liu Y, Zhang Y, Zhao X, Dong A, Du Y, Huang X, Wang J, Lei X, Zheng X. Intravenous transplantation of mesenchymal stem cell improves cardiac performance after acute myocardial ischemia in female rats. Trans. Int. 2006, 19(7): 570–580. DOI:10.1111/j.1432-2277.2006.00307.x
 30. Krishan A, Krishnamurthy H, Totey S. *Applications of flow cytometry in stem cell research and tissue regeneration*. Sci. 2011, 183.
 31. Jaffe BM, Behrman NR. *Methods of hormone radioimmunoassay*. Academic Press 1974.
 32. Santner SJ, Santen RJ, Kulin HE, Demers LM. A model for validation of radioimmunoassay kit reagents: measurement of follitropin and lutropin in blood and urine. Clinical Chemistry. 1981, 27(11): 1892-1895.

33. Nankin HR, Troen P. Repetitive luteinizing hormone elevations in serum of normal men. JCEM. 1971, 33(3): 558-560.
34. Luangpirom A, Kourchampa W, Junaimuang T. Attenuating effect of *Allium ascalonicum* L. on paracetamol induced seminal quality impairment in mice. J. Med. Plan. Res. 2012, 6: 2655–2659. DOI:10.5897/JMPR11.1602.
35. El-Maddawy ZK, El-Sayed YS. Comparative analysis of the protective effects of curcumin and N-acetyl cysteine against paracetamol-induced hepatic, renal, and testicular toxicity in Wistar rats. Environ. Sci. Pollut. Res. Int. 2018, 25 (4): 3468–3479. DOI: 10.1007/s11356-017-0750-3.
36. AbdRabou MA, Mehany ABM, Farrag IM, Belal A, Abdelzaher OF, El-Sharkawy A, Abd El-Azez AM, EL-Sharkawy SM, Al Badawi MH. Therapeutic effect of murine bone marrow-derived mesenchymal stromal/stem cells and human placental extract on testicular toxicity resulting from doxorubicin in rats. Biomed Res Int. 2021, 1-13. DOI:10.1155/2021/9979670.
37. Sabry SA, Sakr SM, Yousef HN. Ameliorating effects of mesenchymal stem cells on testicular injury in a prepubescent male rat model of streptozotocin-induced diabetes. African J. Biol. Sci. 2022, 18 (2): 139-159. DOI: 10.21608/AJBS.2022.174605.1044.
38. Lue Y, Erkkila K, Liu PY, Kimberley Ma, Wang C, Hikim AS, Swerdloff RS. Fate of bone marrow stem cells transplanted into the testis: potential implication for men with testicular failure. Am J Pathol. 2007, 170 (3): 899–908. DOI: 10.2353/ajpath.2007.060543.
39. Akondi BR, Challa SR, Akula A. Protective effects of rutin and naringin in testicular ischemia-reperfusion induced oxidative stress in rats. J Reprod Infertil. 2011, 12(3):209-214.
40. Abarikwu SO, Njoku RC, John IG, Amadi BA, Mgbudom-Okah CJ, Onuah CL. Antioxidant and anti-inflammatory protective effects of rutin and kolaviron against busulfan-induced testicular injuries in rats. Syst Biol Reprod Med. 2022, 68 (2): 151-161. DOI: 10.1080/19396368.2021.1989727.
41. Jahan S, Munawar A, Razak S, Anam S, Ain QU, Ullah H, Afsar T, Abulmeaty M, Almajwal A. Ameliorative effects of rutin against cisplatin-induced reproductive toxicity in male rats. BMC Urol. 2018, 18:107. DOI:10.1186/s12894-018-0421-9.
42. Hassan AI, Alam SS. Evaluation of mesenchymal stem cells in treatment of infertility in male rats. Stem Cell Res. Thera. 2014, 5:131.1-5.