

بنغازي



جامعة

مجلة جامعة بنغازي العلمية



العدد الثاني للمجلد 33 لسنة 2020م

<http://sjuob.uob.edu.ly>

كلمة العدد:

بسم الله الرحمن الرحيم

الحمد لله رب العالمين حمداً كثيراً على نعمته التي أنعم الله بها علينا وألهمنا و وفقنا لإنجاز هذا العمل. والصلاة والسلام على معلم البشرية سيدنا محمد صلى الله عليه وعلى آله وصحبه وسلم تسليماً. إنه لمن دواعي سرورنا أن نشهد بتوفيق من الله صدور العدد الثاني للمجلد 33 من مجلة جامعة بنغازي العلمية والتي مكّنت البَحّاث المهتمين في مجال العلوم الانسانية والتطبيقية والطبية من نشر أبحاثهم على صفحات هذه المجلة لتوفير وإعطاء المعلومة والنتيجة الصحيحة لطلاب العلم والمعرفة، ومن هذا المنطلق فإن هيئة تحرير المجلة تجدد حرصها الدائم على استمرارية صدور المجلة برصانة ومنهجية في البحث العلمي وذلك بإتباع الأساليب العلمية المحكمة في تقييم البحوث العلمية المقدمة من الأساتذة والبَحّاث بإشراف أساتذة متخصصين أخذين في الاعتبار الطرق المتبعة في المجالات العلمية العريقة ، وبهذه المناسبة يسر هيئة التحرير بالمجلة أن تتّمن عالياً جهود جميع البَحّاث والأساتذة المهتمين الذين لم تتنهم الصعاب ولم تؤثر فيهم الظروف الشديدة التي تمر بها بلادنا الحبيبة على تقديم كل ما لديهم من جهد للبحث العلمي واختاروا هذه المجلة لنشر أبحاثهم و أوراقهم العلمية، كما لا يفوتنا أن نتقدم بجزيل الشكر والعرفان لكل من ساهم في تحرير ومراجعة البحوث المقدمة للمجلة وتقديم هذا الانتاج العلمي للوجود.

والله الموفق وعليها التكلان

رئيس التحرير

د. ابتسام عمران العرفي

مجلة جامعة بنغازي العلمية

المجلد الثالث والثلاثون - العدد الثاني

تُعنى بمختلف فروع المعرفة الإنسانية والتطبيقية والطبية

تنشر بحوث باللغتين العربية والإنجليزية

أسرة التحرير

د. ابتسام عمران العرفي	كلية الأسنان	رئيس التحرير
د. عبدالسلام محمد عبدالحفيظ	كلية الاقتصاد	مدير التحرير
د. رجاء محمد بوهادي	كلية القانون	عضواً عن العلوم الانسانية
د. عبير حسين عامر	كلية الطبّ البشري	عضواً عن العلوم الطبية
د. كنز امحمد بوزيد	كلية تقنية المعلومات	عضواً عن العلوم التطبيقية
د. ابتسام علي العبار	كلية الآداب	عضواً عن العلوم الانسانية
أ. د. أحمد محمد الكوافي	كلية العلوم	عضواً عن العلوم التطبيقية
د. مجيد محمد حبريشة	كلية الآداب	عضواً ومدققاً لغوياً
أ. غالية سالم البزار		مقرراً

الموظفون

خليفة أبوشناف الأثرم

للاتصال بنا

مجلة جامعة بنغازي العلمية - جامعة بنغازي مكتب رقم 12 جامعة بنغازي - الحميضة

Email : sjuob@uob.edu.ly

هـ : 0925123074 – 0913823170

شروط النشر في المجلة (Publication's Guidelines):

1. أن يكون النشر في المجلة باللغتين العربية أو الإنجليزية في جميع المجالات التطبيقية والانسانية والطبية.
2. أن يكون البحث العلمي اصيلاً وتتوافر فيه شروط والأصول العلمية والمنهجية المتعارف عليها.
3. أن يتحمل الباحث المسؤولية القانونية عن المعلومات والحقائق الواردة في البحث.
4. ألا يكون البحث قد سبق نشره ورقياً أو إلكترونياً أو مقدم للنشر.
5. يجب أن لا تزيد عدد صفحات البحث عن 25 صفحة.
6. إتباع البحث لنظام (فانكوفر) للتوثيق: [/https://libguides.murdoch.edu.au/Vancouver](https://libguides.murdoch.edu.au/Vancouver).
7. خضوع البحث للتحكيم والموافقة العلمية قبل النشر.
8. تنتقل حقوق الطبع والنشر فور اعتماده .
9. للمجلة الحق في عدم نشر أي بحث يتعارض مع هذه الشروط دون ذكر الأسباب.
10. أن يكون النشر في المجلة إلكترونياً أو ورقياً.

قواعد النشر (Publication's Rules):

1. يجب أن يكتب على صفحة مستقلة عنوان البحث واسم المؤلف (المؤلفون) وعنوانه.
2. يوضع في بداية البحث ملخص باللغتين العربية والإنجليزية، بحيث يتناول الملخص: مشكلة وأهمية البحث، وطبيعة البيانات، وأهم النتائج والتوصيات.
3. يتبع الملخص الكلمات المفتاحية لأهم المصطلحات المستخدمة في البحث.
4. يجب أن تستخدم جميع المواد في كل صفحة حجم صفحة يتوافق مع A4 داخل مستطيل 21×29.7 سم ، يتم توسيطه في الصفحة بدءاً من 1.9 سم من أعلى الصفحة وينتهي بـ 2.5 سم من الأسفل. ويجب أن يكون الهامش الأيمن والأيسر 1.9 سم ، ويجب أن تكون الورقة بتنسيق عمودين 8.20 سم بمساحة 0.80 سم بين الأعمدة.
5. يجب استخدام خط Times New Roman من 9 نقاط ، أو خطوط رومانية أخرى مع محارف معينة .
6. يجب أن يكون العنوان (Times New Roman 16-bold) وأسماء المؤلفين (Times New Roman Regular 11-point) والانتضاءات (Times New Roman 10-point in Italic) ويجب أن يكون عنوان البريد الإلكتروني (Courier Regular 9-point) .
7. بالنسبة للصفحات الأخرى غير الصفحة الأولى ، ابدأ من أعلى الصفحة ، واستمر في تنسيق العمود المزدوج. يجب أن يكون العمودين في الصفحة الأخيرة قريبين من الطول المتساوي قدر الإمكان.

محتويات العدد الثاني للمجلد 33 من مجلة جامعة بنغازي العلمية (2020)

Table of Contents for Volume 33 Issue 2 of the Scientific Journal of University of Benghazi (2020)

م	العنوان	المؤلف	الصفحة
العلوم الإنسانية (Humanities)			
1	لفظة (أزر) في القرآن الكريم	مجيد محمد حبريشة	07
2	واقع الطلاق في المجتمع الليبي	يوسف محمد أبو القاسم الصيد	14
3	واقع استخدام الوسائل التعليمية في التدريس لتلاميذ الشق الثاني من مرحلة التعليم الأساسي العام ببلدية الأبيار	عيسى رمضان محمد مخلوف وعبد الحكيم عبد الحميد بوشنيف موسى و فرج فرجاني محمد عقيلة	24
4	ادارة الازمات الاسرية وعلاقتها بإدارة الصف لدى المعلمات الامهات	إبراهيم أبو عقيل وشهد رفيق نيروخ	38
5	أثر التغيير التنظيمي على عملية اتخاذ القرارات الإدارية	بشير محمد العبار وعادل رابح يونس	50
6	قياس أهم محددات التضخم في الاقتصاد الليبي	على سعيد عبدالله الشريف وعبدالناصر محمد المسلاتي	60
7	دراسة جدوى مشروع إنشاء مستشفى تعليمي استثماري بجامعة عمر المختار	علي محمود فارس وعبد المطلب أحمد بوفروة وحنان عبد الرحمن بركة	70
8	أثر مشاركة العميل على نواتج الخدمات المصرفية المقدمة في المصارف التجارية الليبية العامة العاملة في مدينة بنغازي	فيصل سالم الكيخيا ومريم محمد حسن	81
9	إطار مقترح لتطوير دور الأجهزة الرقابية في تقييم أنظمة الرقابة الداخلية بالمؤسسات الحكومية الليبية	أحمد علي الكاديكي ونرمين خليفة النعاس	96
10	اختبار القدرة التنبؤية للنموذج اللوجستي لـ (Estrella et al , 2000) للتنبؤ بتعثر المصارف	إبراهيم مسعود الفرجاني	108
11	آثار الضرائب على بعض المتغيرات الاقتصادية حسب المنهج الكينزي	جمال نصر الشيباني	116
12	دعم ثقافة الجودة في مؤسسات التعليم العالي: دراسة حالة جامعة بنغازي	عياد طاهر بن إسماعيل	120
13	تحليل مستوى الإفصاح عن المخاطر في التقارير المالية السنوية للمصارف التجارية الليبية	وليد إبراهيم محمد البرغثي ومنيعة حامد حرم سيدهم	130
14	المعوقات التي تحد من التحول نحو الإدارة الإلكترونية في هيئة الرقابة الإدارية دراسة حالة بهيئة الرقابة الإدارية فرع إجدابيا	خالد إقليوان القطراني	143
15	The Impact of Incentives on the Performance Level of Nursing Staff in the Libyan Public Health Sector	Abdelsalam M. A. Saad Fathi S. M. Omar Mohamed M. B. Elfallah	152

محتويات العدد الثاني للمجلد 33 من مجلة جامعة بنغازي العلمية (2020)

Table of Contents for Volume 33 Issue 2 of the Scientific Journal of University of Benghazi (2020)

العلوم التطبيقية (Applied Sciences)			
164	ناصر محمد المسلاتي وسالم هلال الشريف وسميرة عبد الرازق الكواش	دراسة أثر الانفتاح الاقتصادي على النمو الاقتصادي في ليبيا باستخدام نموذج (ARDL) خلال الفترة (1990 - 2014)	1
169	Osama R. Shaltami 1 Fares F. Fares Farag M. EL Oshebi Hwedi Errishi Mohammed S. Aljazwi Abdurabbah. S. Saleh Ali F. Muftah Rachelle R. Favaloro Abla. A. Rhouma	Primarily Assessment of Mamuniyat Formation Sandstones for Glass Industry, Idri Area, SW Libya	2
174	Idress A. Al Gehani Sami A. Alasheebi Ali M. Kalifa Hwedi Errishi Adim A. Hamad	Physiological Responses of Squash (Cucurbita pepo L.) to Humic Acid Treatment under NaCl Stress Conditions	3
180	Osama R. Shaltami Ahmed M. Muftah Moftah, H. El-Shawaihdi Osama A. El-Fallah	Geochemical Characterization Of the U1 Member, Sahabi formation In Northeast Sirt Basin, Libya	4
189	Hanan Salem Abd Alhafid	Existence and unique of the mild solution of stochastic integro differential equation	5
193	Sumaia E. Eshim Amer A. Boushaala Mohammed M.Hamed	Hybrid Genetic Simulated Annealing Algorithm for Job Shop Scheduling Problem	6
العلوم الطبية (Medical Sciences)			
199	Salem Elfleleh Mohammed shultami Halima A.Bargaty Hussain AlKhamry Salem Omaer Salah JABER	Risk of Middle Ear Complications in Intubated Patients on Mechanical Ventilation Versus Non Intubated	1
203	Naeima Betamar Esra Burgeia AbdulGafar Fareg	Evaluation of the Success Rate of Indirect Pulp Capping Treatment using Different Materials: A Clinical Study	2
211	Hoda Mansur Patricia Murray	Investigating The Effect Of Serum Concentration On The Expression Pattern Of Pluripotency And Lineage-Specific Markers In Mouse Embryoid Bodies Using Immunofluorescence	3
220	Rugea Mahmud Mahmmed Namah AL-Sadik EL-Houni	Effect Of Maternal Iron Deficiency Anemia On The Baby Anthropometric Measurements And Hematological Indices	4
226	Gadalla R.M Muftah H.M Aljelane L.K Aldresy A.H Abdualsalm S.F Mohamed N.S Alfrjane H.H1	The Role of Total Quality Management (TQM) in Improving the Health Services Provided at the Benghazi Medical Center (BMC)	5
233	Abdelraouf A. A. Khatal Akram Y. Yasear Laila R. A. Eljreiby Salh S.Muftah	Micromorphological Studies On Implantation Of Albino Rat Endometrium	6

العلوم الطبية

Medical Sciences

Evaluation of the Success Rate of Indirect Pulp Capping Treatment using Different Materials: A Clinical Study

Naeima Betamar^{1*}, Esra Burgeia², AbdulGafar Fareg³

¹ Associate Professor, Department of Conservative Dentistry and Endodontics, Faculty of Dentistry, University of Benghazi, Libya.

² Internship student, Faculty of Dentistry, University of Benghazi.

³ Associate Professor, Department of Statistics, Faculty of Science, University of Benghazi.

Received: 19 / 10 / 2020; Accepted: 31 / 12 / 2020

المخلص:

الهدف من هذه الدراسة السريرية هو تقييم معدل نجاح علاج تغطية اللب غير المباشرة (IPC) باستخدام ثلاث مواد؛ Biner LC و Dycal و TheraCal LC في أسنان الضواك والأضراس الدائمة. في هذه الدراسة تم اختيار عدد 200 من الأسنان الخلفية الدائمة ذات التسوس العميق موصوفة لعلاج اللب غير المباشر (IPC). عدد المرضى 130 مريضاً من كلا الجنسين تتراوح أعمارهم بين 18-55 عاماً. تم تقسيم الأسنان إلى ثلاث مجموعات حسب المادة المستخدمة في تغطية اللب. تم إجراء علاج IPC باستخدام Dycal و Biner LC و TheraCal LC في 67 و 68 و 65 سنناً على التوالي. تم ترميم الأسنان باستخدام مركب راتينج نانو. تم تقييم المرضى سريرياً في 8 أسابيع و 6 أشهر و عام واحد لتقييم نجاح علاج IPC. تم تحليل البيانات إحصائياً باستخدام برنامج SPSS. لوحظ ارتفاع معدل نجاح علاج IPC مع مواد تغطية اللب الثلاثة ولم يكن هناك فرق معتمد به إحصائياً بين المجموعات الثلاث التي تم فحصها ($P>0.05$). في فترة تقييم لمدة عام واحد كان معدل نجاح Dycal و Biner LC و TheraCal LC 97% و 97.1 و 98.5 % على التوالي. أظهرت المواد الثلاثة نسبة نجاح عالية ووجدت أنها ناجحة جداً كمادة IPC ؛ ومع ذلك ، أظهر TheraCal LC أفضل النتائج حتى فترة متابعة لمدة عام واحد. حافظت معالجة I PC على حيوية اللب ووظيفة الأسنان الدائمة، وأن نجاح علاج IPC لا يعتمد على مادة تغطية اللب.

الكلمات المفتاحية:

تسوس عميق، تغطية غير مباشرة لللب، هيدروكسيد الكالسيوم، Dycal، Biner LC، TheraCal LC.

Abstract

The aim of this clinical study was to evaluate the success rate of indirect pulp capping (IPC) treatment with three pulp capping materials; Dycal, Biner LC, and TheraCal LC in permanent premolars and molars teeth.

A total of 200 permanent posterior teeth with deep carious lesions indicated for IPC treatment were included in the study. The number of patients was 130 of both genders, age 18-55 years old. Teeth were divided into three groups according to the material used for pulp capping. IPC treatment was performed using Dycal, Biner LC, and TheraCal LC in 67, 68, and 65 teeth respectively. Teeth were then restored with nanohybrid resin composite. Patients were clinically evaluated at 8 weeks, 6 months, and one year follow up period for the success of IPC treatment. Data were statistically analyzed using SPSS software.

A high success rate of IPC treatment was observed with the three pulp capping materials and that there was no statistically significant difference between the three investigated groups ($P>0.05$). At one year evaluation period, the success rate of Dycal, Biner LC, and TheraCal LC was 97%, 97.1, and 98.5% respectively.

The three materials were exhibited a high success rate and found to be very successful as IPC agents; however, TheraCal LC showed the best results up to one year follow-up period. IPC treatment maintained the pulp vitality and function of the restored permanent teeth, and that the success of IPC treatment is independent of the pulp capping material.

Keywords: Deep caries, Indirect pulp capping, Dycal, Calcium hydroxide, Biner LC, TheraCal LC.

1. INTRODUCTION

The main objective of the modern approach of restorative dentistry is to preserve pulp health of caries teeth, thus reducing the need for root canal treatment and preserve the teeth on the dental arch long-term.¹ Treatment of deep carious lesions is challenging and varied among dental practitioners. The successful treatment is rather a major problem for many of them owing to the complex etiology and pathology of dental caries. Nowadays, based on the biological concept of treatment

strategies along with the development of dental materials; deep caries lesions treated with selective caries removal technique which is recommended by a recent report of European Society of Endodontology.²⁻⁴

Indirect pulp capping (IPC) is a therapeutic intervention in the treatment of deep carious lesions approximating the pulp of vital teeth with no signs of pulp degeneration, to avoid pulp exposure and preserve pulp vitality.^{1,5-7} IPC is generally used in deep cavity preparation, involves the removal of all soft,

*Correspondence:

Naeima Betamar
Faculty of Science, University of Benghazi, Libya.
Nabetamar@gmail.com

demineralized infected dentine with or without leaving behind a layer of caries-affected dentine remains in close to the pulp, followed by application of pulp capping material, and a restoration^{5,8,9}. The idea behind this technique based on the fact that dental pulp can develop inorganic mineralized dentine-like matrix as a part of the repair mechanism in the dentine pulp complex.^{7, 10} i.e. preserving the pulp biologic function to stimulate and induce odontoblasts to produce tertiary dentine. The objective of IPC treatment is to discontinue demineralization of carious dentine and thus arresting the carious progression by promoting remineralization and stimulating reparative dentine formation.¹⁰

Several materials have been used and tested as pulp capping agents.¹⁰⁻¹⁶ Important properties of these materials are biocompatibility, preserve pulp vitality, and antibacterial property.¹⁷ Moreover, other properties being able to promote tissue repair, adhere to dentine and restorative material, as well as sealing capabilities.¹⁸ Among those materials calcium hydroxide Ca(OH)₂ which was introduced by Hermann in 1921 has been considered the "gold standard" of direct and indirect pulp capping for several decades.^{16, 17} It is available in several forms such as aqueous suspensions, cements, liners, or filled resins as visible light-cured liner containing calcium hydroxide.¹⁷ For instance; Dycal (Dentsply, Milford, DE, USA) is a Ca(OH)₂ based material, a self-cure radiopaque liner with an alkaline pH (pH 9–11).¹⁹ In clinical practice, it is the first material applied as a liner in patients with deep cavities and remains the most popular material among dentists.^{19, 20}

Ca(OH)₂-based materials are reported to display excellent antibacterial properties and induce mineralization and reparative dentine formation because of their chemical irritation nature.¹⁶ However, the conventional two-paste chemical-cure formulation of Ca(OH)₂ cement has several disadvantages that may fail of the treatment such as high solubility in oral fluids that lead to dissolution over time, lack of adhesion to dentine as well as to resin-based restorative materials, poor sealing ability, internal resorption, extensive dentine formation obliterating the pulp chamber, low elastic modulus, and low compressive strength.^{8, 17, 19, 21} Furthermore, degradation after acid etching and presence of tunnel defects through reparative dentine bridge.^{8, 16, 19} Due to these drawbacks of chemical-cure Ca(OH)₂; a light-cured,

single component liner contains calcium hydroxide and is polymerized by visible light was introduced in 1988 with improved strength, no solubility in acid, and minimal solubility in water.²²

Various other materials have been introduced into a dental practice and are used as pulp capping agents such as glass ionomer, resin-modified glass ionomer cements and adhesives, and calcium-silicate materials (CSMs).^{2, 16} CSMs are recent bioactive materials that can induce regenerative responses in the human body and form hard tissue dentine bridge.^{13, 16} These materials are superior in biocompatibility, and sealing of the pulp capped site, and result in more predictable clinical outcomes.^{13, 16} An example of these materials is mineral trioxide aggregate (MTA),¹⁶ that has gained an excellent reputation and commonly recommended and preferred by dentists.²³ The second generation of these CSMs are; Biodentine and TheraCal LC.^{1, 2} Biodentine is new bioactive tricalcium silicate-based cement, designed to be used as a permanent, biocompatible dentine substitute and stimulates pulp cells to form tertiary dentine.² TheraCal LC is a light-cured flowable resin-modified calcium silicate material introduced by Bisco Inc. Schaumburg, IL, USA in 2011, designed to overcome the weaknesses of the previous generation such as poor bonding of CSMs to resin restorations. It is used as direct and indirect pulp capping material that enables the immediate application of final restoration.^{1, 2} TheraCal bonds to deep moist dentine, and performs as a barrier and protectant material of the pulp-dentine complex underneath composite, amalgam, and cement.²⁴ Hence, it acts as a suitable replacement for Ca(OH)₂, and several other pulp capping materials.¹⁶ TheraCal displayed strong physical properties, high radiopacity, and lower solubility than either ProRoot MTA or Dycal.¹ The aim of this clinical study was to evaluate the success rate of IPC treatment using three materials namely; Dycal, Binar LC, and TheraCal LC in permanent posterior teeth with deep carious lesions.

2. MATERIALS AND METHODS

Detailed descriptions and composition of materials used in the study are listed in Table 1.

Table 1: Description and composition of the materials used in the study

Material	Components*	Manufacturer
Dycal® Ivory	Base: disalicylate ester, butylene glycol; calcium phosphate; calcium tungstate; zinc oxide; iron oxide. Catalyst: calcium hydroxide; ethylene toluenesulfonamide; zinc stearate; titanium dioxide; zinc oxide; iron oxide.	Dentsply
Binar LC	UDMA resin, calcium dihydroxide, dimethylaminoethyl-methacrylate, barium aluminium silicate, TEGDMA. photoinitiator.	Voco GmbH, Cuxhaven, Germany
TheraCal LC	CaO, calcium silicate particles (Portland cement type III), Sr glass, fumed silica, barium sulphate, barium zirconate, Bis-GMA, HEMA, and PEGDMA.	Bisco Inc., Schamburg, IL, USA
Filtek™ Z250 XT	Zirconia/silica 20 nm, fillers loading 82% by weight. Bis-GMA, UDMA, Bis-EMA, TEGDMA, and PEGDMA.	3M ESPE
Adper™ Single Bond 2	10% by weight 5nm colloidal silica nanofiller. Bis-GMA, ethanol, water, photoinitiator, polyacrylic and polyitaconic acids, HEMA.	3M ESPE

*According to manufacturers' technique profiles: Bis-GMA: Bisphenol-A-glycidyl dimethacrylate; UDMA: Urethane dimethacrylate; Bis-EMA: Bisphenol A ethoxylate dimethacrylate; TEGDMA: Triethylene glycol dimethacrylate; PEGDMA: Polyethylene glycol dimethacrylate; HEMA; 2-hydroxyethyl methacrylate. CaO: calcium oxide

Study design:

A randomized clinical trial was conducted at Al-Raja Dental Clinic in Benghazi City. A total number of 200 posterior permanent teeth with deep carious lesion indicated for indirect pulp capping treatment from 130 adult patients were selected in this study. Informed consent from the patients was obtained after giving a brief explanation of the kind of investigation and the clinical procedure that was to be conducted. Inclusion criteria included both genders, age 18 to 55 years old having premolars or molars teeth with a deep carious lesion on the occlusal or occluso-proximal surface with no pain or with very mild sensitivity to cold or discomfort when chewing. Preoperative radiograph shows deep carious lesion extending greater than two-third of dentine thickness approaching pulp,¹¹ but with a definite radiodense region between the deepest part of caries lesion and the pulp.²⁵ Teeth with clinical symptoms of irreversible pulpitis, a history of prolonged spontaneous pulpal pain were excluded. In addition, teeth with pulp necrosis or with a negative response to pulp tests, mobility, tender on percussion, presence of swelling, or fistula were also excluded. Furthermore radiographically; if teeth show the presence of periapical or furcation area radiolucency,¹¹ periapical pathologies, internal or external root resorption, absence of a normal appearance of the periodontal ligament, were also excluded from the study.

Clinical procedures:

A thermal vitality test was done to testify tooth sensitivity; after removal of the stimulus, the response rapidly disappeared. A preoperative periapical radiograph was taken for each patient to assess penetration depth and the extent of the caries lesion, as well as the thickness of the remaining dentine overlying the pulp chamber. All the clinical procedures were performed by the standard method of IPC treatment under a rubber dam. After clinical examination and radiographic assessment, local anesthesia was administered. A sterile diamond bur suitable to the size of the cavity in a high-speed handpiece was used for cavity preparation under constant water cooling. Caries tissue was removed completely from the lateral walls and cavosurface margins of the cavity preparation. Excavation of carious dentine included removal of all active demineralized carious tissue from the peripheral sites to the center or the deepest part of the lesion. Not to mention that it is usually at the pulpal floor or the axial wall in the case of the class II cavity. Selective caries removal was performed with the cautious excavation of soft, wet, and necrotic caries tissue reaching to a non-soft, relatively firm dentine, accompanied by visual inspection and tactile sensation to avoid pulp exposure. A hand instrument spoon excavator was used to remove necrotic fragments of caries tissue followed by a round bur in a low-speed handpiece under water coolant. A new sterile bur is replaced when approaching the deeper area of the caries lesion, to reduce the number of microorganisms in the cavity and to avoid the introduction of infected dentine into the pulp which might lead to irreversible inflammation and treatment failure.^{26, 27} The cavity was then washed with normal saline and gently dried. Teeth were then randomly assigned into three groups; Gp:1, Gp:2, and Gp:3 according to the material used for pulp capping. Gp1 consists of 67 teeth allocated to receive treatment with Dycal (chemical-cured radiopaque $\text{Ca}(\text{OH})_2$). Dycal is available as a two-paste system; a catalyst paste and a base paste. It was mixed with equal quantities of both the catalyst and the base to a homogenous paste and applied with a dycal applicator directly over the deepest spots of

the dentine on the pulpal floor of the cavity preparation. The excess dycal material was removed from the surrounding cavity walls and enamel margins. Gp 2 consists of 68 teeth and treated with Biner LC (Light-curing radiopaque one-component cavity liner containing $\text{Ca}(\text{OH})_2$). It was applied to the deepest spots of the pulpal floor and then light-cured according to manufacturer instructions. TheraCal LC (Light-curing, resin-modified calcium silicate filled liner) was applied in Gp3 similarly as of Gp2, and consists of 65 teeth. Biner LC and TheraCal LC are supplied by the manufacturer in syringes and disposable tips and requiring no preparation before use. Excess material was removed while still soft and before light curing. The randomization procedure was performed as follows; a number corresponding to each IPC treatment group was written on a piece of paper and kept in a closed box. A paper was selected from the box by a person other than the operator, and the treatment indicated was carried out.⁹ The bonding and restorative procedures were performed likewise for the three investigating groups as followed; The cavity walls and margins were acid etched with 37% phosphoric acid semi gel (Meta Biomed Co Ltd., Korea) according to the manufacturer's instructions, then thoroughly rinsed off with water and gently air-dried using compressed air without desiccation. Bonding agent Adper™ Single Bond 2 Adhesive (3M ESPE) was applied with a microbrush in two consecutive layers. A gentle stream of air was applied from an air syringe to allow evaporation of the solvent then light-cured for 10 seconds with LED light-curing unit (Mini LED, Satelec, France). For class II cavity preparation, a Tofflirmire retainer with a matrix band was placed before acid etching and bonding procedures. Nanohybrid resin composite restorative material Filtek™ Z250 XT (3M ESPE) was incrementally packed in the cavity preparation. Each increment was polymerized for 20s using the same LED light-curing unit. The occlusal adjustment was then performed in maximum intercuspation and eccentric movements. The identified high spots were carefully removed using extra-fine grit diamond burs EX-17EF, FO-23EF (Toboom Shanghai Precise Abrasive Tool Co., Ltd) with water coolant, and then polished with polishing tips to eliminate any surfaces scratches (Enhance Dentsply Caulk). After completing the treatment, the patient was instructed about preventive measures and maintenance of oral hygiene.

At follow-up evaluation, a detailed clinical examination was performed for every patient at 8 weeks, 6 months, and one year intervals. Periapical radiographs were taken at the end of the clinical trial. However, the patient who returned to the dental clinic with pain or any sign of treatment failure, a periapical radiograph was taken for him/her to evaluate the restoration and periapical region that could be attributed to a failure of the treatment.²⁰

The clinical success of the treatment was evaluated by the following criteria: intact restoration, positive (normal) response to cold pulp test, absence of prolonged spontaneous pain, no tenderness on percussion or palpation, no tooth mobility, no abscess or swelling of periodontal tissues of the treated tooth.^{9, 14, 28}

On radiographic examination; the success of the treatment included intact lamina dura, no periapical radiolucency, no internal or external root resorption.^{14, 28, 9, 20} Any tooth presented with one of the above-mentioned criteria was recorded as treatment failure and was root canal treated, or replacement of the restoration when indicated. One tooth was restored at each clinical visit. Restorative treatments, clinical

and radiographic evaluations were performed by the same operator.¹⁴ The data were statistically analyzed using SPSS software (version 16, SPSS Inc. IBM Corp. Chicago, USA). Chi-Square test was applied and the level of significance was set as $P < 0.05$.

3. RESULTS

At the beginning of the study, 213 teeth received IPC treatment. By the end of the one year, 13 patients were lost to follow up. Hence a total of 200 experimental teeth were evaluated. The mean age was 30.7 ± 8.4 yrs. The numbers of female and male patients were 102 (78.5%) and 28 (21.5%) respectively. From all teeth included in the study, 150 (75%) were molars and 50 (25%) were premolars. Among those, the upper right first molar and lower left second molar received the highest number of restorations; 23 (11.5%) and 22 (11%) respectively (Figure 1). Additionally, 70 (35.5%) restorations were class I, and 129 (64.5%) were class II. A total of 67 (33.5%) teeth treated with Dycal, 68 (34%) teeth treated with Biner LC, and 65 (32.5%) teeth received TheraCal LC.

Statistical analysis revealed no significant differences in the success rate of IPC between the three investigated groups using three pulp capping materials ($P > 0.05$) (Table 2). Clinical and radiographic evaluation revealed that the success rate of IPC treatment at one year evaluation period was 97% for Dycal (GP1), where 56 teeth considered successful and only 2 teeth (3.0%) were failed. These two teeth are UR4 presented with pain during the 8-weeks evaluation period and went for root canal treatment (RCT), and UL6 attended at a one year evaluation period with fractured restoration. The success rate for Biner LC (Gp2) was 97.1%; where 66 teeth were considered successful. Two teeth (2.9%) recorded failure; UL5 and LL6 and hence root canal treated. Whereas for TheraCal (Gp3); the success rate was 98.5%. Sixty-four teeth treated were considered successful and only one tooth (1.5%), UL5 reported failure, and went for RCT. In general, all the failed teeth belonged to class II cavity preparations, in which three of them were premolars and two teeth were molars. Radiographic examination at the end of one year evaluation period exhibited intact lamina dura, absence of periapical radiolucency, no internal or external root resorption.




Figure 1: The number and frequency of all restored teeth with the three pulp capping materials

Table 2: Results of the success and failure of the three IPC materials				
Pulp capping materials (Group)	N	Success	Failure	P-value
Dycal (Gp1)	67	65 97.0%	2 3.0%	0.833
Biner LC (Gp2)	68	66 97.1%	2 2.9%	
TheraCal (Gp3)	65	64 98.5%	1 1.5%	
Total	200	195 97.5%	5 2.5%	

4. DISCUSSION

Management of deep caries lesion is an area of continuous amendment and challenge to the dentists, in terms of which treatment will produce the best outcomes and excellent prognosis which is the preservation of pulp vitality of the tooth with no apical pathology.²⁹ Deep caries can be recognized by two layers: infected dentine which is a superficial layer of soft demineralized destructive dentine tissues infiltrated with a large number of bacteria and denatured collagen fibrils that cannot be remineralized; and the other layer is the affected dentine, which is a deeper layer of partially demineralized dentine with intact collagen fibrils that can be remineralized.^{30, 31} It is hard to differentiate exactly these two layers in the clinical reality. However, using some indicators such as the color and consistency of the decayed dentine as well as the operator skills and clinical experience to make the good decision for excavation of the infected dentine, leaving only the affected dentine to avoid pulp exposure.^{32, 33} Yet, if the pulp happens to be exposed and infected during caries removal, a successful outcome will be reduced.²⁹ Therefore, a conservative procedure has been suggested to avoid pulp exposure by using the indirect pulp capping (IPC) treatment, which can be either a one-step or two-steps clinical procedure.³⁴ It has been suggested to apply a cavity liner and/or base if caries extends close to the pulpal tissue^{7, 30, 31} i.e. when the remaining dentine thickness (RDT) is ≤ 0.5 mm, as a protective layer under the final restorations and to coat the innermost layer of caries-affected dentine to induce remineralization.^{7, 12, 20} In the current clinical trial a one-step (one-visit) IPC treatment under controlled isolation was performed. The choice for IPC was based on a careful pulp diagnosis with proper evaluation of the pain history, symptoms, clinical and radiographic findings.¹⁵ In addition, selective and cautious caries removal was done with expectations that reparative dentine would be formed, demineralized dentine would be remineralized, and the number of viable bacteria would be reduced, thus caries lesion is arrested.³ Evidence has been shown that the two clinical visits IPC treatment would not be necessary, because, through the second visit, the treatment procedure increased the risks of pulp exposure.³⁵ Additionally, some patients do not attend the second clinical visit which might result in partial or total loss of the temporary filling and therefore failure of the treatment.⁹ This would add extra cost, time, and discomfort to the patient, and would thus be

detrimental to pulpal health^{9, 33, 35, 36} The IPC treatment for each patient was firstly evaluated after 8 weeks, based on the fact that the evaluation period of 8- to 12- weeks post-treatment is considered sufficient period to promote remineralization of carious dentine and pulp recovering^{26, 37} In their study, the investigators found a statistically significant increase in total mineral content in the testing samples with a qualitative increase in radiopacity of the calcium hydroxide²⁶

In the current study, few cases reported failures among the three investigated groups; Dycal, Biner LC, and TheraCal LC treated teeth. The main reason for the failure could be attributed to a non-diagnostic inflammation in the pulp that could not be precisely diagnosed before starting the treatment. On several occasions, dentists find difficulty in evaluating the true clinical status of the pulpal tissue underneath deep caries lesion, which sometimes makes it hard to reach a precise and accurate diagnosis of pulpal health. In this context, it has been documented that the correct and accurate “gold standard” of pulp health status is the histological analysis, and cannot be determined by clinical signs, symptoms, or radiologic appearance⁸ Instead, Bjørndal³⁸ in 2008 reported that there was no reliable method to determine the degree of pulp inflammation in deep carious lesions, which is the main reason why successful treatment of deep caries is still uncertain. It is worth mentioning that all failure teeth belonged to class II cavity preparations. The multi-surface restorations such as class II (DO, MO, and MOD) produced inferior results compared with one-surface restorations such as Class I occlusal cavity, which is established in a previous study.³⁹

Regarding the dycal treated teeth; it has been documented that the older $\text{Ca}(\text{OH})_2$ formulations do not provide a long-term biological seal, protection, and permanent barrier against bacterial infection and microleakage. Because $\text{Ca}(\text{OH})_2$ cement dissolve within 1–2 years, and tunnel defects in the majority of the reparative dentin bridges underneath the capping material.^{2, 19} However, according to Accorinte et al.,⁴⁰ by the time the $\text{Ca}(\text{OH})_2$ is vanished due to dissolution, dentine bridge has formed. Yet, in the present study up to a one-year follow-up period dycal treated teeth (Gp1) exhibited a high success rate (97%). Dycal has been widely used and commonly preferred in clinical practice as a pulp capping material with the longest track record of clinical success.⁸ The high success rate observed with dycal treated teeth could be attributed to its effect on pulp repair by one or more of several mechanisms of action.⁸ It is an alkaline material, biocompatible, induces remineralization, and reduces the risk of bacterial infection.^{2, 41} In addition to the ability of $\text{Ca}(\text{OH})_2$ to release hydroxyl (OH) and calcium (Ca) ions upon its dissociation into the surrounding environment.^{17, 42} These ions caused a chemical injury that stimulates the pulp to perform a defense mechanism where the undifferentiated cells within the pulp distinguish into odontoblasts that form a hard mineralized tissue as a reparative dentine.⁴² According to Murray et al.,⁴³ $\text{Ca}(\text{OH})_2$ liners mediate and maintain the health and vitality of the underlying odontoblasts survival for the deposition of reparative dentine when the RDT is ≤ 0.5 mm. Furthermore, the hydroxyl ions diffuse into the dentine and create an alkaline pH level, which is an unsuitable environment for remaining bacteria in the cavity, therefore penetration of bacteria into the pulp is minimized.^{8, 18, 21} In this context, the high pH of $\text{Ca}(\text{OH})_2$ destroy the bacterial cell membrane and protein structure, and thus sufficiently terminate the residual

bacteria.^{17, 19} This may explain the high success rate and the few numbers of failures among this investigating group, which is in agreement with other studies who reported a high success rate with $\text{Ca}(\text{OH})_2$ ranging from 70% to 91%.^{9, 20}

Biner LC treated teeth (Gp2) also exhibited a high success rate (97.1%). There were only two teeth reported failures (2.9%) soon after treatment and therefore root canal treated (RCT). Biner LC is a visible light-cured (VLC) liner flow on-demand, easy handling, and dispensing, consists of calcium hydroxide and barium sulfate dispersed in a urethane dimethacrylate resin containing initiators and accelerators activated by visible light.²² The polymeric resins of this material permit a good bond with the overlying composite restoration, and also reported to be compatible with adhesive systems, dentine, and pulp.^{21, 22} However, Gandolfi et al.,¹³ found that the amount of calcium released by VLC liner was very small compared to that released by the dycal. Nevertheless, other studies reported that the results of pulp capping with different adhesive resin-based systems are nearly equal to those with $\text{Ca}(\text{OH})_2$ in dentine bridge formation.⁴⁴ It has been suggested to use VLC $\text{Ca}(\text{OH})_2$ liner as indirect pulp capping material underneath all types of restorations, and if a dentine bonding agent is preferred to be used, the VLC liner should only be applied on the deepest dentine leaving the rest of the cavity surface free for bonding.¹⁷

Only one tooth (1.5%) reported failure with TheraCal LC treated teeth, which consider a very high success rate (98.5%) of the treatment. This material is recently available in the market, designed for use in indirect and direct pulp capping. The main advantages of this material are the high biocompatibility, bioactivity property, and intrinsic osteoconductive activity which is the ability to induce a biological and regenerative response in the human body, resulting in the formation of a bond between the material and the tissue which is the dentine bridge of improved quality and superior sealing ability of the pulp capped site.^{2, 13, 24} Additionally, TheraCal LC consists of tricalcium silicate particles in a hydrophilic monomer that displayed higher and significant calcium release and lower solubility than Dycal making it a stable and durable material as a liner or base.^{24, 32} These good properties could clarify the excellent results obtained with TheraCal treated teeth in this study. On the other hand, this material is “white” in color and opaque hence it should be placed in a thin layer so as not to show through composite materials and affecting the shade of restoration.¹⁶

Since this material has been recently presented to the dental practitioners and researchers, only a few clinical studies were evaluating its performance, and the majority of them were investigated primary teeth, or when TheraCal was used as direct pulp capping. Clinical studies have shown that TheraCal LC can encourage reparative dentine and dentine bridge formation when used as direct and indirect pulp capping material.^{45, 46} Another study reported that all experimental teeth (20) treated with TheraCal LC remained vital after 2 years.⁴⁷ Petrolo et al.,⁴⁸ found that most vital teeth were seen in the TheraCal group after a 2 years evaluation period when they investigated the effects of TheraCal on pulp with a self-etching adhesive system.

It is worth mentioning that very few teeth reported failures with resin-contained pulp capping materials; i.e. Biner LC and TheraCal LC. Literature reported that the components of the adhesive resin-based liners are found to be toxic in cell cultures

which may threaten the pulp in the short-term,⁴⁹ in addition to the temperature rise during photopolymerization of these materials.⁵⁰ It has been documented that the presence of resin in the components of pulp capping material which may remain unpolymerized is often accompanying by adverse pulpal reactions that lead to pulp toxicity and inflammation even in the absence of bacteria.^{49, 51} Thus inflamed pulp due to caries would have decreased healing capability.⁴⁹ Moreover, TheraCal could induce adverse pulpal effects when applied in pulp capping procedures despite the low heat generation of the photopolymerization process.⁵⁰ These are the possible explanations for the failures that occurred with the resin-contained pulp capping materials though they were very few cases.

A high success rate of IPC treatment was observed among the three experimental groups regardless of the material used throughout the study. This success can be principally related to several factors such as; the careful selection of the cases, proper pulpal diagnosis, adequate and proper excavation of caries infected dentine, good isolation, and prevention of contamination. In addition to a good choice of IPC medication as well as the restorative material used for the permanent restoration.^{32, 52} The good quality bond between the restorative material and tooth structure which in turn provide an effective seal of the lesion from the oral environment is very important to prevent bacterial substrate infiltrating into the dentine and to control the progression of caries.^{8, 52} Studies have demonstrated a significant decrease in the count of bacterial under sealed restorations.^{41, 53} In addition, other studies have reported that; clinical, radiographic, and microbiological evaluations have demonstrated that sealing the cavity properly could lead to arrest the deep active carious lesions even when inert materials such as wax or gutta-percha were used as liners.^{54, 55}

Finally, it is sensible to consider that pulp capping material individually did not influence the clinical success of treatment for deep caries lesions of permanent teeth, which is also specified in the Cochrane review that indicated that the difference in base materials did not show any changes.²⁹ It's a combination of several factors as mentioned formerly in the discussion section which is in agreement with other studies.^{3, 7, 20, 32} In addition to the correct diagnosis of the health status of the pulp before starting the treatment and maintain the pulp vitality after treatment.

5. CONCLUSIONS:

Within the limitations of this study, the following conclusions can be drawn: IPC treatment maintained the pulp vitality and function of the restored permanent teeth, and that the success of IPC treatment seen in the present study is independent of the pulp capping material. Proper and careful case selection included healthy pulp and correct pulpal diagnosis, good isolation, a good marginal seal of the cavity preparation with a suitable restoration are important factors for successful treatment. All three materials were exhibited a high success rate and found to be very successful as IPC agents, however, TheraCal LC seems promising material and exhibited the best results up to a one year follow-up period. Additional research including controlled clinical studies with larger sample size and longer-term follow up evaluation periods is required to better assess and understand the clinical performance and efficiency of these relatively new materials in the long run.

6. REFERENCES

1. Zaparde N, Gunda S, Patil A. TheraCal... Future of Pulp Capping??? . International J of Development Research 2017;7(10):16338-42.
2. Kunert M, Lukomska-Szymanska M. Bio-Inductive Materials in Direct and Indirect Pulp Capping—A Review Article. J Materials 2020;13, 1204:1-20.
3. Schwendicke F, Frencken JE, Bjørndal L, et al. Managing Carious Lesions: Consensus Recommendations on Carious Tissue Removal Adv Dent Res 2016a;28:58–67.
4. Duncan HF, Galler KM, Tomson PL, et al. European Society of Endodontology position statement: Management of deep caries and the exposed pulp. Int Endod J 2019;52:923–34.
5. Costa CA, Hebling J, Hanks CT. Current status of pulp capping with dentin adhesive systems: A review. Dent Mater 2000;16:188–97.
6. Thompson V, Craig RG, Curro FA, Green WS, Ship JA. Treatment of deep carious lesions by complete excavation or partial removal: A critical review. J Am Dent Assoc 2008;139(6):705-12.
7. Schwendicke F, Gøestemeyer G, Gluud C. Cavity lining after excavating caries lesions: meta-analysis and trial sequential analysis of randomized clinical trials. Journal of Dentistry 2015;43:1291–7.
8. Hilton TJ. Keys to clinical success with pulp capping: a review of the literature. Oper Dent 2009;34(5):615-25.
9. Maltz M, Garcia F, Garcia R, et al. Randomized Trial of Partial vs. Stepwise Caries Removal: 3-year Follow-up. J Dent Res 2012;91(11):1026-31.
10. Cao CY, Mei ML, Li QL, Lo ECM, Chu CH. Methods for biomimetic remineralization of human dentine: A systematic review. Int J Mol Sci 2015;16:4615–27.
11. Gruythuysen RJ, van Strijp AJ, Wu MK. Long-term survival of indirect pulp treatment performed in primary and permanent teeth with clinically diagnosed deep carious lesions. J Endod 2010;36:1490-3.
12. Renata F, Luciano C, Alice SP, et al. Clinical and radiographic evaluation of indirect pulp treatment in primary molars: 36 months follow-up. Am J Dent 2007;20(3):189-92.
13. Gandolfi MG, Siboni F, Botero T, et al. Calcium silicate and calcium hydroxide materials for pulp capping: Biointeractivity, porosity, solubility and bioactivity of current formulations. J Appl Biomater Funct Mater 2015;13(1):41–60.
14. Gurcan AT, Seymen F. Clinical and radiographic evaluation of indirect pulp capping with three different materials: a 2-year follow-up study. European Journal of Paediatric Dentistry 2019;20(2):105-10.
15. Lucia CD, Raluca D, Levente GC. Clinical and radiographic evaluation of indirect pulp capping with MTA and calcium hydroxide for permanent teeth. Romanian Journal of Medical and Dental Education 2019;8(3):35-45.
16. Qureshi A, Soujanya E, kumar N, kumar P, Shivarao S. Recent Advances in Pulp Capping Materials: An Overview. Journal of Clinical and Diagnostic Research 2014;8(1):316-21.

17. Arandi ZN. Calcium hydroxide liners: a literature review. *Clinical, Cosmetic and Investigational Dentistry* 2017; 9:67–72.
18. Mantellini MG, Botero TM, Yaman P, et al. Adhesive resin induces apoptosis and cell cycle arrest of pulp cells. *Journal of Dental Research* 2003;82(8):592–96.
19. Cox CF, Subay RK, Ostro E, Suzuki S, Suzuki SH. Tunnel defects in dentin bridges: Their formation following direct pulp capping. *Oper Dent* 1996;21(1):4-11.
20. da Rosa WLO, Lima VP, Moraes RR, Piva E, da Silva AF. Is a calcium hydroxide liner necessary in the treatment of deep caries lesions? A systematic review and meta-analysis. *International Endodontic J* 2019;52(5):588–603.
21. Schuurs AH, Gruythuysen RJ, Wesselinde PR. Pulp capping with Resin based composite versus calcium hydroxide: A review endodontics. *Dental Trauma* 2000;16:240-50.
22. Sakaguchi RL, Powers JM. *Craig's Restorative Dental Materials*. Philadelphia, PA: Elsevier/Mosby; 2012.
23. Dominguez MS, Witherspoon DE, Gutmann JL, Opperman LA. Histological and scanning electron microscopy assessment of various vital pulp-therapy materials. *Journal of Endodontics* 2003;29(5):324-33.
24. Gandolfi MG, Siboni F, Prati C. Chemical-physical properties of TheraCal, a novel light-curable MTA-like material for pulp capping. *Int Endod J* 2012;45(6):571-9.
25. Bjørndal L, Fransson H, Bruun G, et al. Randomized Clinical Trials on Deep Carious Lesions: 5-Year Follow-up. *J Dent Res* 2017;96(7):747-53.
26. Carlos AC. Remineralization of Carious Dentine: In Vitro Microradiographic Study in Human Teeth Capped by Calcium Hydroxide. *Braz Dent J* 2004;15:59-62.
27. Dammaschke T, Galler K, Krastl G. Current recommendations for vital pulp treatment. *Dtsch Zahnärztl Z Int* 2019;1:43-52.
28. Venkatesh Babu NS, Kavyashree BS, Amitha HA. Comparative Assessment of Success Rate of Indirect Pulp Treatment With 2% Chlorhexidine Gluconate Disinfecting Solution, Calcium Hydroxide And Resin Modified Glass Ionomer Liner In Primary Teeth - A Prospective Study. *Journal of Dental and Medical Sciences* 2018;17(1):68-77.
29. Miyashita H, Worthington HV, Qualtrough A, Plasschaert A. Pulp management for caries in adults: maintaining pulp vitality. *Cochrane Database Systematic Review* 2007;18(2):CD004484
30. Heymann H, Swift EJ, Ritter AV, Sturdevant CM, editors. *Sturdevant's Art and Science of Operative Dentistry*. 6th ed. St Louis: Elsevier/ Mosby; 2013.
31. Hilton TJ, Ferracane JL, Broome JC, editors. *Summitt's Fundamentals of Operative Dentistry: A Contemporary Approach*. 4th ed. Hanover Park: Quintessence Publishing Co; 2013.
32. Alex G. Direct and Indirect Pulp Capping: A Brief History, Material Innovations, and Clinical Case Report. *Compendium* 2018;39(3):182-88.
33. Bjørndal L, Reit C, Bruun G, al. e. Treatment of deep caries lesions in adults: randomized clinical trials comparing stepwise vs. direct complete excavation, and direct pulp capping vs. partial pulpotomy. *Eur J Oral Sci* 2010;118(3):290-97.
34. Bjørndal L. Dentin and pulp reactions to caries and operative treatment: biological variables affecting treatment outcome. *Endod Topics* 2002;2(1):10-23.
35. Kidd EA. How “clean” must a cavity be before restoration? . *Caries Res* 2004;38(3):305–13.
36. Ricketts D, Innes Nb, Schwendicke F. *Selective Removal of Carious Tissue Monogr Oral Sci*. Basel, Karger 2018;27:82-91.
37. Franzon R, Casagrande L, Pinto AS, et al. Clinical and radiographic evaluation of indirect pulp treatment in primary molars: 36 months follow-up. *Am J Dent* 2007;20(3):189-92.
38. Bjørndal L. Indirect pulp therapy and stepwise excavation. *J Endod* 2008;34(7):Suppl: S29-S33.
39. Maltz M, Alves LS, Jardim JJ, Moura Mdos S, De Oliveira EF. Incomplete caries removal in deep lesions: a 10-year prospective study. *Am J Dent* 2011;24:211–14
40. Accorinte M, Reis A, Loguerio A, de Araújo V, Muench A. Influence of rubber dam isolation on human pulp responses after capping with calcium hydroxide and an adhesive system. *Quintessence International* 2006;37(3):205–12.
41. Maltz M, Oliveira EF, Fontanella V, Bianchi R. A clinical, microbiologic, and radiographic study of deep caries lesions after incomplete caries removal. *Quintessence Int* 2002;33(2):151–59.
42. Fulzele P, Baliga S, Thosar N, Pradhan D. Evaluation of calcium ion, hydroxyl ion release and pH levels in various calcium hydroxide based intracanal medicaments: an in vitro study. *Contemporary Clinical Dentistry* 2011;2(4):291– 95.
43. Murray PE, Smith AJ, Windsor LJ, Mjor IA. Remaining dentine thickness and human pulp responses. *Int Endod J* 2003;36(1):33–43.
44. Akimoto N, Momoi Y, Konoh A, et al. Biocompatibility of Clearfil Liner Bond 2 and Clearfil AP-X system on nonexposed and exposed primate teeth. *Quint Int* 1998;29(3):177–88.
45. Cannon M, Gerodias N, Viera A, Percinoto C, Jurado R. Primate pulpal healing after exposure and TheraCal application. *J Clin Pediatric Dent* 2014;38(4):333-37.
46. Menon NP, Varma BR, Janardhanan S, et al. Clinical and radiographic comparison of indirect pulp capping treatment using primary molars: a randomized clinical trial. *Contemp clin Dent* 2016;7(4):475-80.
47. Nicola S, Petrolo F, Allegra C, et al. 2-years clinical outcome of direct pulp capping with resin-based materials. *J Dent Res* 2015;94(spec iss A). Abstract 3709.
48. Petrolo F, Comba A, Scansetti M, et al. Effects of light-cured MTA like material on direct pulp capping. . *Dental Materials* 2014;Jan 1;30:e151.

49. Hebling J, Lessa FCR, Nogueira I, de Carvalho RM, de Costa CAS. Cytotoxicity of resin based light-cured liners The American Journal of Dentistry 2009;22(3):137–42.
50. Savas S, Botsali MS, Kucukyilmaz E, Sari T. Evaluation of temperature changes in the pulp chamber during polymerization of light-cured pulp-capping materials by using a VALO LED light curing unit at different curing distances. Dent. Mater. 2014;33:764–69.
51. Jeanneau C, Laurent P, Rombouts C, Giraud T, About I. Light-cured Tricalcium Silicate Toxicity to the Dental Pulp. J. Endod. 2017;43:2074–80.
52. Pereira MA, Santos-Junior RB, Tavares JA, et al. No additional benefit of using a calcium hydroxide liner during stepwise caries removal: a randomized clinical trial. J Amer Dent Associ 2017;148(6):369–76.
53. Maltz M, Oliveira E, Fontanella V, Carminatti G. Deep caries lesions after incomplete dentine caries removal: 40-month follow-up study. Caries Research 2007;41(6):493–96.
54. De Oliveira EF, Carminatti G, Fontanella V, Maltz M. The monitoring of deep caries lesions after incomplete dentine caries removal: results after 14-18 months. Clinical Oral Investigation 2006;10:134–9.
55. Pinto AS, Araújo FB, Franzon R, et al. Clinical and microbiological effect of calcium hydroxide protection in indirect pulp capping in primary teeth. Am J Dent 2006;19:382–86.

University Of Benghazi



Scientific Journal of University of Benghazi



Volume33,Issue 2,2020

<http://sjuob.uob.edu.ly>