

## Diagnostic and Epidemiologic Features of Visceral Leishmaniasis Transmission by The Phlebotomus Fly in Children Admitted to The Children's Hospital in Benghazi

Dareen E. A. Jadullah<sup>1</sup>, Samira S. Garbou<sup>2\*</sup>, Lubna A. Salem<sup>3</sup>, Sara A. Elwarrad<sup>4</sup>

<sup>1</sup> Faculty of Pharmacy, University of Benghazi, Libya.

<sup>2</sup> Department of infectious disease, Faculty of Public Health, University of Benghazi, Benghazi, Libya.

<sup>3</sup> Department of Hematology, Children's Hospital, Benghazi, Libya.

<sup>4</sup> Department of Statistics, Faculty of Arts and Sciences, University of Benghazi, Al-Marg, Libya.

Received: 03 / 12 / 2020; Accepted: 27 / 03 / 2021

### المخلص:

يعتبر داء الليشمانيا عدوى طفيلية تنتقل عادةً بين الحيوانات والإنسان عن طريق لدغة أنثى ذباب رمل الماص للدم، وقد أجريت هذه الدراسة لتسليط الضوء على مرض الليشمانيا الحشوي في الأطفال. حيث يُعد ظهور هذا المرض مصدر قلق خوفاً من انتشاره بصورة كبيرة، وهو مستوطن في مناطق معينة في ليبيا. استند تشخيص داء الليشمانيا الحشوي على تاريخ المريض والتشخيص السريري، وتم تأكيده بأخذ خزعة من السائل الشوكي لكشف الشكل السوطي للطفيل (Amastigotes)، وتُفحص مجهرياً بعد صبغ المسحة بصبغة جيمسا (Giemsa) حيث تعتبر هذه الطريقة نهج الخط الأول في تشخيص مرض الليشمانيا.

بالإضافة إلى دراسة مقطعية مستعرضة أجريت في الفترة من 1 يناير 2013 إلى 31 ديسمبر 2019 للتحقيق في العوامل الوبائية لمرض داء الليشمانيا الحشوي لدى الأطفال الذين تم إدخالهم إلى مستشفى الأطفال في بنغازي، ليبيا.

أشارت النتائج إلى أنه خلال فترة الدراسة كان هناك 51 طفل مصاب بمرض الليشمانيا الحشوي من بين 99 حالة اشتباه. وكانت الأعراض السريرية الأكثر هي نقص الكريات الحمر وتضخم النوى وتضخم الكبد والطحال والحمى المطولة. ونوع الليشمانيا الذي يسبب المرض هي ليشمانيا دونوفانيا (Leishmania Donovan) وقد وجدنا الفئة العمرية من 0 - 3 سنوات هم الأكثر تضرراً ما بين الأطفال، وظهرت نتائج الدراسة زيادة إصابة الأطفال الذكور مقارنةً بالاطفال الإناث 2 : 1، جغرافياً أعلى إصابة كانت في مدينة مرزق الريفية 5.45٪. وقد خلصت الدراسة إلى أن داء الليشمانيا يشكل خطراً صحياً ووبائياً في عدة أماكن في ليبيا حيث يصيب الأطفال والبالغين. وسيكون من المهم أن نفهم بعناية كيفية انتقال المرض بين الأشخاص، وزيادة الوعي المهني والمجتمعي، وتحسين التشخيص، وتوفير طرق العلاج المناسبة.

### الكلمات المفتاحية:

داء الليشمانيا الحشوي، الأطفال، مستشفى الأطفال، بنغازي، ليبيا.

### Abstract

The Leishmania species are the causative agents of leishmaniasis, which is typically transmitted between vertebrate hosts through the bite of blood-sucking female phlebotomine sand flies. This study was conducted due to the paucity of information in the literature relevant to visceral leishmaniasis (VL) in Libyan children. The emergence of visceral leishmaniasis is of concern in Libya and is endemic in certain regions. Diagnosis of visceral leishmaniasis was based on patient history, physical results, and then confirmed by laboratory bone marrow examination. A cross-sectional, hospital-based study was conducted from the 1st of January 2013 to the 31st of December 2019 to investigate the epidemiological factors of VL in children admitted to the Children Hospital in Benghazi, Libya. The results indicated that during the study period there were 51 children diagnosed with VL among 99 suspected cases. The most common clinical symptoms were myelopoiesis (90.9%), aspiration (90.9%), cellularity (89.9%), erythrocytopenia (88.9%), megakaryopoiesis (88.9%), hepatosplenomegaly (58.6%) and prolonged fever (48.5%). Children in the age group  $\geq 3$  years were most affected, and the male-female ratio was 2: 1. The highest number of cases of children were from the rural city of Murzuq at 45.5%. The study concluded that VL is an endemic health threat affecting children and adults in Libya. Considering that a portion of locally identified Leishmania donovani strains possess the ability to vascularize, it will be important to carefully understand the characteristics of transmission, increase professional and community awareness as well as improving prognosis and the availability of appropriate methods of treatment.

**Keywords:** visceral leishmaniasis, children, the Children's Hospital, Benghazi, Libya.

## 1. INTRODUCTION

Visceral leishmaniasis (VL), or kala-azar "Black fever", is a vector-borne disease and the most serious form of leishmaniasis. It is caused by the protozoan parasites Leishmania donovani and L. infantum. VL is transmitted to humans through the bite of infected female sand flies, genus phlebotomus [1]. In the past

decade, the number of leishmaniasis cases has increased, both in endemic countries and all over the world [2]. Recent estimates showed about 50 000 to 90 000 new cases of VL. Only about 25–45% of cases which occur each year worldwide are reported to the WHO [3]. This increase may be associated with the increased frequency of travel, ecotourism activity, military operations and immigration, according to an Italian study [4].

\*Correspondence:

Samira S. Garbou. Department of infectious disease, Faculty of Public Health, University of Benghazi, Benghazi, Libya.

[samira.garboui@gmail.com](mailto:samira.garboui@gmail.com)

Since VL is a fatal disease when left untreated, and because treatments used have high toxicity, a very sensitive and specific diagnostic test is required. The gold method for diagnosing VL is fundamentally the demonstration of the parasite by microscopic examination of splenic aspirate culture. Spleen aspiration can be associated with hemorrhage, and the operation should only be done with access to surgical facilities. For this reason, the most common clinical samples used for this purpose are bone marrow and lymph node aspirates for parasitological diagnosis. The specificity of these methods is high, but the sensitivity differs depending on the type of specimen [5, 6, 7]. In Libya, VL was identified from north-eastern Libya and southern Saharan and sub-Saharan areas where many of the reported VL patients had traveled outside of Libya and the majority were young children [8, 9]. In these areas, there are two endemic foci of infantile visceral leishmaniasis (IVL), caused by the pathogen *L. infantum*, MON-1. [10]. IVL infects children under 5 years, in Al-Jabel Al-khaddar (northeastern region) and Ubari (southern region) [8, 9, 11]. It is characterized by irregular bouts of fever, weight loss, enlargement of the spleen and liver, and anaemia [2].

The exact prevalence of VL in Libya was unknown since most scientific research focused on cutaneous leishmaniasis in terms of the epidemic [12], Spatiotemporal analysis [13] and vector surveillance [14] compared to visceral leishmaniasis, where few papers dealt with the diagnosis, epidemiology, and treatment of the VL disease.

Therefore, this study aimed to shed light on the diagnosis used to recognize the disease in Benghazi's Children's Hospital, where many cases who were less than 15 years old came from various eastern and southern cities.

As well as knowing the spread of the epidemic and the number of cases during the five-year study period. In addition to the demographic characteristics of all cases, the clinical evaluation of patients and the treatments that are given and follow-up cases were investigated.

## 2. MATERIAL AND METHODS

### 2.1. Patients and samples

Clinical specimens were obtained from 99 patients, clinically diagnosed with visceral leishmaniasis admitted to the Children's Hospital in Benghazi, the main hospital and reference for the treatment of VL and other diseases in northeastern Libya. Between 2013 and 2019, clinical features and laboratory findings related to children were examined.

Permission for conducting this study was obtained by the Human Resources Department in the Children's Hospital administration, which is responsible for research and studies conducted within the hospital.

### 2.2. Patients and epidemiological information

Demographic data for the children was recorded, such as age, gender, nationality, usual residence, medical history and diagnostic year, along with the assumed country/region in which leishmaniasis was contracted. Laboratory finding such as an increase in erythrocyte sedimentation rate (ESR), alanine transaminase (SGPT), and aspartate transaminase (SGOT) and major clinical manifestations such as fever, splenomegaly, hepatomegaly and treatment type.

### 2.3. Diagnosis of VL disease in children

The diagnostic criteria was the detection of a parasite (amastigotes form) in a Giemsa stained bone marrow smear which is a very specific method and the first-line approach in VL diagnosis [15]. Approximately 0.3 ml of bone marrow was aspirated and each bone marrow sample was processed immediately. The samples were not mixed with anti-coagulant to avoid morphological artifacts. The sample was then air-dried, fixed in methanol, and stained with routine stains such as Giemsa or Leishman stain. The diagnosis is considered confirmed if at least one laboratory test is conducted.

### 2.4. Statistical analysis

The data was analyzed using SPSS Version 21 and Excel 2013

## 3. RESULTS AND DISCUSSION

The confirmation of cases of VL in Benghazi's Children's Hospital took place mainly by laboratory criteria. Table 1 shows the number of cases during the study period and the number of laboratory-confirmed cases. Among the 99 patients that had a diagnosis of VL in their clinical records, bone marrow examination showed that *Leishmania* cells were found in only 51 of these patients. A direct examination of the bone marrow shows that all positive cases of VL were infected with the *L. donovani* parasite (Table 1). It is fairly well established that VL in North Africa and the Middle East is caused by *L. donovani* and *L. infantum*. In Libya, *L. donovani* was identified as the causative agent of VL a long time ago [16] and this fact was confirmed by our findings.

The most common clinical symptoms of all hospitalized cases and clinically diagnosed VL were myelopoiesis 90.9%, aspiration 90.9%, followed by cellularity 89.9%, erythropoiesis 88.9%, megakaryopoiesis 88.9%, hepatosplenomegaly 58.6% and prolonged fever 48.5% whereas the anemia percentage was only 4% of children figure 1. About half of these cases correspond to Murzuq in southwestern Libya, which had 45.5% confirmed cases in that period (Figure 2).

Regarding the age group, the interval with the highest incidence of cases was  $\geq 3$  years (51%), followed by 4 to 7 years (30%) while the lowest infection rate of VL was observed in the

age group between 12 to 15 years. In addition, there are several records that have no information (5%). Based on (Table 2) and (Figure 3), the male to female ratio was 1:2 indicating that there was a statistically significant gender difference. Of the total cases reported, 92.9% were Libyan citizens (Table 2). There was an increase in the number of cases that were cured (42.4%) whereas 7.1% died and 50.5% were negative as shown in Table 2.

The number of VL reports had significant growth over the years analyzed, as shown in Figure 4. There were a few cases observed between 2013 and 2014, and in 2016, a significant increase was recorded, followed by a slight decrease in 2017. The reason for the increase in infection cases may be due to the cessation of the control program that was implemented by the National Center for Disease Control, which was an existing and organized program. This program included reducing rodents and the use of vector control methods by local pest management companies. It was able to eliminate 85-90% of desert mice (*Psammomys*), which is the animal reservoir in the target areas [17]. After 2011, the role of the National Center became purely technical, providing technical advice to the municipalities where the disease is endemic. Stopping the control program of the transmitters of the disease

meant an increase in their population, and consequently an increase in the infection in the endemic areas, and its spread to new places where the disease had not previously been present.

It is important to note that the decline in the number of cases to zero in 2015 does not mean that there were no incidences of the disease, as in this year the city of Benghazi was in a state of war, and all the cases from the south, which is known to have the highest rate of the disease went to Tripoli for treatment.

The analysis of the monthly distribution for all years of the study of cases of visceral leishmaniasis showed that there are periods with a greater prevalence of the disease. Generally, between October and January is where the highest peaks of the disease are concentrated, especially in December followed by January, as shown in Figure 5.

IVL is known to occur in North Africa [18]. In Libya, there are two endemic foci of (IVL), the northern coastal areas near Tripoli and Al-Jabel Al-khdar (northeastern region) [10]. Most infected cases are under 5 years old. A Tunisian study including 1089 cases revealed that about 50% of VL cases were aged between 1 and 3 years [19], which confirms what we discovered in this study. The sex ratio was male biased as males remain near the animal host for a longer time than females do, however, the difference is not significant. This data is in accordance with previous findings [20], IVL infects children under 5 years, in Al-Jabel Al-khdar and Ubary (Southern region) [8, 16]. The southern area has reclaimed land for agriculture with increasing water resources and some other environmental changes, which may be partly responsible for creating these new foci [9]. The disease is endemic in South Libya, and rates have been rising due to the country's prolonged crisis. The current study revealed that the majority of cases of children come from the city of Murzuq; similar finding were observed by various authors [21, 22, 23]. This may be due to complex emergencies and mass population movements, in addition to what was mentioned above about the cessation of control programs in the country.

#### 4. CONCLUSION

Parasitological diagnosis with a demonstration of the amastigote forms in tissues remains the only method of diagnosis used in the hospital. Epidemiologically, it was observed that Murzuq has a considerable percentage of children with VL, which highlights the need for strategies to reduce the disease. After concluding that the vast majority of infected children are from the Southern areas, there is a need for a wider healthcare system so that the patient does not need to travel to get the appropriate treatment, as it is essential that therapy be carried out until the healing process is complete.

There also must be better means of disseminating information so that people can be aware of what the recommendations for prevention and control of leishmaniasis are. The Environmental Protection Agency must also continuously control infected dogs and thus prevent the transmission of the disease to humans.

The data analyzed in this paper regarding VL in the Children's Hospital in Benghazi cannot be extended to cover the whole country, since it depends on the patient's presence in the hospital from various cities and some patients may go to hospitals in the capital, Tripoli. Thus, we point out the limitations of this research.

#### 5. RECOMMENDATIONS



Ideally, the combination of multiple methods is recommended for leishmaniasis diagnosis such as serologic detection of antibodies to recombinant K39 antigen. In addition, Polymerase chain reaction (PCR) is a common molecular technique that is highly sensitive and specific. The control of VL disease depends on focusing on the most common areas, while treatments depend on several factors such as the patient's characteristics (e.g. age, immune status, and kidney/liver function), the extent of disease, availability of drugs, associated infections and previous treatments.


**Table 1:** Method of identification, causative Leishmania species and study period.

| Leishmania spp. | Study period | Suspected cases by clinical examination | Confirmed cases by bone marrow examination | Total number of confirmed cases |
|-----------------|--------------|-----------------------------------------|--------------------------------------------|---------------------------------|
| L. Donovan      | 2013         | 1                                       | 1                                          | 51                              |
| "               | 2014         | 2                                       | 2                                          |                                 |
| "               | 2015         | 0                                       | 0                                          |                                 |
| "               | 2016         | 10                                      | 6                                          |                                 |
| "               | 2017         | 19                                      | 10                                         |                                 |
| "               | 2018         | 32                                      | 17                                         |                                 |
| "               | 2019         | 35                                      | 15                                         |                                 |


**Table 2:** Demographic variables of cases admitted to the Children's Hospital

| Demographic Variables | Positive | Negative | Total |
|-----------------------|----------|----------|-------|
| <b>Age groups</b>     |          |          |       |
| Miss                  | 1        | 4        | 5     |
| ≥3                    | 23       | 28       | 51    |
| 4-7                   | 17       | 12       | 29    |
| 8-11                  | 7        | 2        | 9     |
| 12-15                 | 3        | 2        | 5     |
| <b>Gender</b>         |          |          |       |
| Male                  | 33       | 28       | 61    |
| Female                | 18       | 20       | 38    |
| <b>Nationality</b>    |          |          |       |
| Libyan                | 49       | 43       | 92    |
| Non Libyan            | 2        | 5        | 7     |
| <b>Mortality</b>      |          |          |       |
| Unfollow              | 2        | 48       | 50    |
| Treated               | 42       | 0        | 42    |
| Dead                  | 7        | 0        | 7     |
| Total                 | 51       | 48       | 99    |


**Figure 1.** The frequency of clinical symptoms in children admitted with VL to the Children's Hospital between 2013 and 2019.**Figure 2:** Geographical distribution of children admitted with VL to the Children's Hospital between 2013 and 2019.



**Figure 3:** Number of IVL district by gender during the period of 2013-2019



**Figure 4:** Incidence rate and number of children's cases of VL from 2013 to 2019



**Figure 5:** Distribution of confirmed cases of VL from January to December in the years between 2013 and 2019.

## 6. REFERENCES

- Levin, M L, 2014. Medical entomology for students. *Emerging infectious diseases*, 20(8), p.1428.
- World Health Organization. Report on the interregional meeting on leishmaniasis among neighbouring endemic countries in the Eastern Mediterranean, African and European regions, Amman, Jordan 23–25 September 2018. No. WHO-EM/CTD/081/E. World Health Organization, Regional Office for the Eastern Mediterranean, 2019.
- World health organization (WHO). Leishmaniasis. Report of WHO 14 March 2019. <https://www.who.int/news-room/fact-sheets/detail/leishmaniasis>
- Di Muccio T, Scalone A, Bruno A, Marangi M, Grande R, Armignacco O, Gradoni L, Gramiccia M. Epidemiology of imported leishmaniasis in Italy: Implications for a European endemic country. *PLoS One*. 2015 Jun 26;10(6):e0129418.
- Gupta AK, Singh A, Srivastava S, Shankar P, Singh S. Visceral Leishmaniasis in Children: Diagnosis, Treatment, and Prevention. *Journal of Pediatric Infectious Diseases*. 2017 Dec;12(04):214-21.
- Antinori S, Calattini S, Longhi E, Bestetti G, Piolini R, Magni C, Orlando G, Gramiccia M, Acquaviva V, Foschi A, Corvasce S. Clinical use of polymerase chain reaction performed on peripheral blood and bone marrow samples for the diagnosis and monitoring of visceral leishmaniasis in HIV-infected and HIV-uninfected patients: a single-center, 8-year experience in Italy and review of the literature. *Clinical infectious diseases*. 2007 Jun 15;44(12):1602-10.
- Chappuis F, Sundar S, Hailu A, Ghalib H, Rijal S, Peeling RW, Alvar J, Boelaert M. Visceral leishmaniasis: what are the needs for diagnosis, treatment and control?. *Nature reviews microbiology*. 2007 Nov;5(11):S7-16.
- Jain S, El Mangoush MA, El-Bouri K, Mahfouz MO. Kala azar in an adult Libyan and review of visceral leishmaniasis in Libya. *Tropical and geographical medicine*. 1990 Jul 1;42(3):283-5.
- Mehabresh MI. Visceral leishmaniasis: new foci of infection in Libya. *The Journal of Tropical Medicine and Hygiene*. 1994 Oct 1;97(5):282-5.
- Gramiccia M. The identification and variability of the parasites causing leishmaniasis in HIV-positive patients in Italy. *Annals of Tropical Medicine & Parasitology*. 2003 Oct 2;97(sup1):65-73.
- Dar FK, Elmauhoub M, Kidwai SA, Mahabresh MI, Saba A, El Bouri KW. Preliminary observations on the epidemiology of visceral leishmaniasis in the Libyan Arab Jamahiriya. *Paediatric Infectious Diseases in Arab Countries*, Elzouki, AY (Ed.). John Wiley & Sons Ltd. 1987:205-10.
- Husain RA, Shaari H, Solla M, Ebrahim HA. Semulation of Leishmaniasis Epidemiology in Libya Using Agent Based Modelling. In 2019 19th International Conference on Sciences and Techniques of Automatic Control and Computer Engineering (STA) 2019 Mar 24 (pp. 324-329). IEEE.
- Amro A, Al-Dwibe H, Gashout A, Moskalenko O, Galafin M, Hamarsheh O, Frohme M, Jaeschke A, Schönián G, Kuhls K. Spatiotemporal and molecular epidemiology of cutaneous leishmaniasis in Libya. *PLoS neglected tropical diseases*. 2017 Sep 7;11(9):e0005873.
- Dokhan MR, Jaouadi K, Salem S, Zenbil O, Gonzalez JP, Salah AB, Annajar BB. Natural Infection of *Phlebotomus sergenti* by *Leishmania tropica* in Libya. *The American journal of tropical medicine and hygiene*. 2018 May 9;98(5):1339-42.
- Garcia LS. *Diagnostic medical parasitology*. American Society for Microbiology Press; 2006 Nov 29.
- Mehabresh MI, El-Mauhoub MM. Visceral leishmaniasis in Libya—review of 21 cases. *Annals of tropical paediatrics*. 1992 Jan 1;12(2):159-63.
- World Health Organization) WHO). Cutaneous leishmaniasis in Libya. WHO 2021. <http://www.emro.who.int/neglected-tropical-diseases/countries/cl-libya.html>
- Aoun K, Ben Abda I, Habboul Z, Lemrani M, Harrat Z, Bouratbine A. Visceral leishmaniasis in North african countries. *PUJ*. 2013;6(1):35-8.
- Aoun K, Jeddi F, Amri F, Ghrab J, Bouratbine A. Actualités épidémiologiques de la leishmaniose viscérale en Tunisie. *Médecine et maladies infectieuses*. 2009 Oct 1;39(10):775-9.
- Fahmy AR, Samy AM, Doha SA, Shehata MG. Preliminary field investigations on *Phlebotomine* sandflies (Diptera: Psychodidae) from a recent cutaneous leishmaniasis focus in Northern-Sinai, Egypt. *Egyptian Academic Journal of Biological Sciences. A, Entomology*. 2009 Jun 1;2(1):9-5.
- Younis EZ, Khater HF, Hayam Elawamy A, Aldinali A, Rabia El Ghazal B. Biochemical Assays and Epidemiological Status of Visceral Leishmaniasis among Patient Attending to Benghazi Children's Hospital. *Ann Microbiol Immunol*. 2020; 3 (1):1022.
- World Health Organization) WHO) LIBYA ANNUAL REPORT 2019 – Humanitarian Response. World Health Organization; 2019. [https://www.humanitarianresponse.info/sites/www.humanitarianresponse.info/files/documents/files/who\\_libya\\_annual\\_report\\_2019.pdf](https://www.humanitarianresponse.info/sites/www.humanitarianresponse.info/files/documents/files/who_libya_annual_report_2019.pdf)
- Kimutai A, Ngure P, Tonui W, Gicheru M, Nyamwamu L. Leishmaniasis in Northern and Western Africa: a review. *African Journal of Infectious Diseases*. 2009;3(1).